

MARCAINE- bupivacaine hydrochloride injection, solution
MARCAINE WITH EPINEPHRINE- bupivacaine hydrochloride and epinephrine bitartrate injection, solution
Hospira, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MARCAINE™ and MARCAINE™ WITH EPINEPHRINE safely and effectively. See full prescribing information for MARCAINE™ and MARCAINE™ WITH EPINEPHRINE.

MARCAINE™ (bupivacaine hydrochloride) injection, for infiltration, perineural, caudal, epidural, or retrobulbar use
MARCAINE™ WITH EPINEPHRINE (bupivacaine hydrochloride and epinephrine) injection, for infiltration, perineural, caudal, or epidural use
Initial U.S. Approval: 1972

WARNING: RISK OF CARDIAC ARREST WITH USE OF MARCAINE IN OBSTETRICAL ANESTHESIA

See full prescribing information for complete boxed warning.

There have been reports of cardiac arrest with difficult resuscitation or death during use of MARCAINE for epidural anesthesia in obstetrical patients. In most cases, this has followed use of the 0.75% (7.5 mg/mL) concentration. Resuscitation has been difficult or impossible despite apparently adequate preparation and appropriate management. Cardiac arrest has occurred after convulsions resulting from systemic toxicity, presumably following unintentional intravascular injection. The 0.75% (7.5 mg/mL) concentration of MARCAINE is not recommended for obstetrical anesthesia and should be reserved for surgical procedures where a high degree of muscle relaxation and prolonged effect are necessary (5.1).

----- **INDICATIONS AND USAGE** -----

MARCAINE contains bupivacaine, an amide local anesthetic, and MARCAINE WITH EPINEPHRINE is a combination of bupivacaine, an amide local anesthetic, and epinephrine, an alpha and beta-adrenergic agonist. MARCAINE / MARCAINE WITH EPINEPHRINE is indicated in adults for the production of local or regional anesthesia or analgesia for surgery, dental and oral surgery procedures, diagnostic and therapeutic procedures, and for obstetrical procedures. For each type of block indicated to produce local or regional anesthesia or analgesia, specific concentrations and presentations are recommended. (1, 2.2)

Limitations of Use

Not all blocks are indicated for use with MARCAINE / MARCAINE WITH EPINEPHRINE given clinically significant risks associated with use. (1, 2.2, 4, 5.1, 5.4, 5.5, 5.7, 5.9)

----- **DOSAGE AND ADMINISTRATION** -----

- Not for intrathecal use. (2.1)
- Avoid use of solutions containing antimicrobial preservatives (i.e., multiple-dose vials) for epidural or caudal anesthesia. (2.1, 5.4)
- Three mL of MARCAINE WITH EPINEPHRINE without antimicrobial preservative (0.5% bupivacaine with 1:200,000 epinephrine) is recommended for use as a test dose prior to caudal and lumbar epidural blocks when clinical conditions permit. (2.4)
- See full prescribing information for:
 - Recommended concentrations and dosages of MARCAINE / MARCAINE WITH EPINEPHRINE according to type of block. (2.2)
 - Additional dosage and administration information pertaining to use in epidural anesthesia, test dose for caudal and lumbar epidural blocks, use in dentistry, and use in ophthalmic surgery. (2.3, 2.4, 2.5, 2.6)

----- **DOSAGE FORMS AND STRENGTHS** -----

MARCAINE injection and MARCAINE WITH EPINEPHRINE injection are available in multiple concentrations. See full prescribing information for detailed description of each formulation. (3)

CONTRAINDICATIONS

- Obstetrical paracervical block anesthesia. Its use in this technique has resulted in fetal bradycardia and death. (4)
- Intravenous regional anesthesia (Bier Block). (4)
- Known hypersensitivity to bupivacaine or to any local anesthetic agent of the amide-type or to other components of MARCAINE / MARCAINE WITH EPINEPHRINE. (4)

WARNINGS AND PRECAUTIONS

- Dose-Related Toxicity: Monitor cardiovascular and respiratory vital signs and patient's state of consciousness after injection of MARCAINE / MARCAINE WITH EPINEPHRINE. (5.2)
- Methemoglobinemia: Cases of methemoglobinemia have been reported in association with local anesthetic use. See full prescribing information for more detail on managing these risks. (5.3)
- Chondrolysis with Intra-Articular Infusion: Intra-articular infusions of local anesthetics including MARCAINE following arthroscopic and other surgical procedures is an unapproved use, and there have been post-marketing reports of chondrolysis in patients receiving such infusions. (5.5)
- Risk of Cardiac Arrest with Intravenous Regional Anesthesia Use (Bier Block): There have been reports of cardiac arrest and death during the use of bupivacaine for intravenous regional anesthesia (Bier Block). (5.7)
- Allergic-Type Reactions to Sulfites in MARCAINE WITH EPINEPHRINE: MARCAINE WITH EPINEPHRINE contains sodium metabisulfite, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. (5.8)
- Risk of Systemic Toxicities with Unintended Intravascular or Intrathecal Injection: Unintended intravascular or intrathecal injection may be associated with systemic toxicities, including CNS or cardiorespiratory depression and coma, progressing ultimately to respiratory arrest. Aspirate for blood or cerebrospinal fluid (where applicable) prior to each dose and consider using a test dose of MARCAINE WITH EPINEPHRINE. (5.9)

ADVERSE REACTIONS

Most common adverse reactions are related to the central nervous system and the cardiovascular system. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Local Anesthetics: The toxic effects of local anesthetics are additive. Monitor for neurologic and cardiovascular effects when additional local anesthetics are administered. (7.1)
- Monoamine Oxidase Inhibitors and Tricyclic Antidepressants: Administration of MARCAINE WITH EPINEPHRINE to patients receiving monoamine oxidase inhibitors or tricyclic antidepressants may produce severe, prolonged hypertension. Concurrent use of these agents should generally be avoided. (5.6, 7.2)
- Ergot-Type Oxytocic Drugs: Concurrent administration of MARCAINE WITH EPINEPHRINE and ergot-type oxytocic drugs may cause severe, persistent hypertension or cerebrovascular accidents. (5.6, 7.3)
- Nonselective Beta-Adrenergic Antagonists: Administration of MARCAINE WITH EPINEPHRINE (containing a vasoconstrictor) in patients receiving nonselective beta-adrenergic antagonists may cause severe hypertension and bradycardia. Concurrent use of these agents should generally be avoided. (5.6, 7.4)
- Drugs Associated with Methemoglobinemia: Patients are at increased risk of developing methemoglobinemia when concurrently exposed to nitrates, nitrites, local anesthetics, antineoplastic agents, antibiotics, antimalarials, anticonvulsants, and other drugs. (7.5)
- Potent Inhalation Anesthetics: Serious dose-related cardiac arrhythmias may occur if preparations containing a vasoconstrictor such as epinephrine are used in patients during or following the administration of potent inhalation anesthetics. (5.13, 7.6)

USE IN SPECIFIC POPULATIONS

- Pediatric Use: Administration of MARCAINE / MARCAINE WITH EPINEPHRINE in pediatric patients younger than 12 years is not recommended. (8.4)
- Geriatric Use: Patients 65 years and over, particularly those with hypertension, may be at increased risk for developing hypotension while undergoing anesthesia with MARCAINE / MARCAINE WITH EPINEPHRINE. (8.5)
- Moderate to Severe Hepatic Impairment: Consider increased monitoring for bupivacaine systemic

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 7/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF CARDIAC ARREST WITH USE OF MARCAINE IN OBSTETRICAL ANESTHESIA

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosage and Administration Information
- 2.2 Recommended Concentrations and Dosages of MARCAINE / MARCAINE WITH EPINEPHRINE
- 2.3 Use in Epidural Anesthesia
- 2.4 Test Dose for Caudal and Lumbar Epidural Blocks
- 2.5 Use in Dentistry
- 2.6 Use in Ophthalmic Surgery

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Risk of Cardiac Arrest with Use of MARCAINE in Obstetrical Anesthesia
- 5.2 Dose-Related Toxicity
- 5.3 Methemoglobinemia
- 5.4 Antimicrobial Preservatives in Multiple-Dose Vials
- 5.5 Chondrolysis with Intra-Articular Infusion
- 5.6 Risk of Adverse Reactions Due to Drug Interactions with MARCAINE WITH EPINEPHRINE
- 5.7 Risk of Cardiac Arrest with Intravenous Regional Anesthesia Use (Bier Block)
- 5.8 Allergic-Type Reactions to Sulfites in MARCAINE WITH EPINEPHRINE
- 5.9 Risk of Systemic Toxicities with Unintended Intravascular or Intrathecal Injection
- 5.10 Risk of Toxicity in Patients with Hepatic Impairment
- 5.11 Risk of Use in Patients with Impaired Cardiovascular Function
- 5.12 Risk of Ischemic Injury or Necrosis in Body Areas with Limited Blood Supply
- 5.13 Risk of Cardiac Arrhythmias with Concomitant Use of Potent Inhalation Anesthetics
- 5.14 Risk of Adverse Reactions with Use in Head and Neck Area
- 5.15 Risk of Respiratory Arrest with Use in Ophthalmic Surgery
- 5.16 Risk of Inadvertent Trauma to Tongue, Lips, and Buccal Mucosa in Dental Applications

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- 7.1 Local Anesthetics
- 7.2 Monoamine Oxidase Inhibitors and Tricyclic Antidepressants
- 7.3 Ergot-Type Oxytocic Drugs
- 7.4 Nonselective Beta-Adrenergic Antagonists
- 7.5 Drugs Associated with Methemoglobinemia
- 7.6 Potent Inhalation Anesthetics

7.7 Phenothiazines and Butyrophenones

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF CARDIAC ARREST WITH USE OF MARCAINE IN OBSTETRICAL ANESTHESIA

There have been reports of cardiac arrest with difficult resuscitation or death during use of MARCAINE for epidural anesthesia in obstetrical patients. In most cases, this has followed use of the 0.75% (7.5 mg/mL) concentration. Resuscitation has been difficult or impossible despite apparently adequate preparation and appropriate management. Cardiac arrest has occurred after convulsions resulting from systemic toxicity, presumably following unintentional intravascular injection. The 0.75% (7.5 mg/mL) concentration of MARCAINE is not recommended for obstetrical anesthesia and should be reserved for surgical procedures where a high degree of muscle relaxation and prolonged effect are necessary [see *Warnings and Precautions (5.1)*].

1 INDICATIONS AND USAGE

MARCAINE / MARCAINE WITH EPINEPHRINE is indicated in adults for the production of local or regional anesthesia or analgesia for surgery, dental and oral surgery procedures, diagnostic and therapeutic procedures, and for obstetrical procedures. Specific concentrations and presentations of MARCAINE / MARCAINE WITH EPINEPHRINE are recommended for each type of block indicated to produce local or regional anesthesia or analgesia [see *Dosage and Administration (2.2)*].

Limitations of Use

Not all blocks are indicated for use with MARCAINE / MARCAINE WITH EPINEPHRINE given clinically significant risks associated with use [see *Dosage and Administration (2.2)*, *Contraindications (4)*, *Warnings and Precautions (5.1, 5.4, 5.5, 5.7, 5.9)*].

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

- MARCAINE / MARCAINE WITH EPINEPHRINE is not for intrathecal use.
- Avoid use of MARCAINE / MARCAINE WITH EPINEPHRINE solutions containing antimicrobial preservatives (i.e., multiple-dose vials) for epidural or caudal anesthesia [see *Warnings and Precautions (5.4)*].
- Discard unused portions of solution not containing preservatives, i.e., those supplied in single-dose vials, following initial use.
- Visually inspect this product for particulate matter and discoloration prior to administration whenever solution and container permit. MARCAINE / MARCAINE WITH EPINEPHRINE are clear, colorless solutions. Do not administer solutions which are discolored or contain particulate matter.
- Mixing or the prior or intercurrent use of any other local anesthetic with MARCAINE / MARCAINE WITH EPINEPHRINE is not recommended because of insufficient data on the clinical use of such mixtures.

Administration Precautions

- MARCAINE / MARCAINE WITH EPINEPHRINE are to be administered in carefully adjusted dosages by or under the supervision of experienced clinicians who are well versed in the diagnosis and management of dose-related toxicity and other acute emergencies which might arise from the block to be employed.
- Use MARCAINE / MARCAINE WITH EPINEPHRINE only if the following are immediately available: oxygen, cardiopulmonary resuscitative equipment and drugs, and the personnel resources needed for proper management of toxic reactions and related emergencies [see *Warnings and Precautions (5.2)*, *Adverse Reactions (6)*, *Overdosage (10)*].
- The toxic effects of local anesthetics are additive. Monitor for neurologic and cardiovascular effects related to local anesthetic systemic toxicity when additional local anesthetics are administered with MARCAINE / MARCAINE WITH EPINEPHRINE [see *Warnings and Precautions (5.2)*, *Drug Interactions (7.1)*, *Overdosage (10)*].
- Aspirate for blood or cerebrospinal fluid (where applicable) prior to injecting MARCAINE / MARCAINE WITH EPINEPHRINE, both the initial dose and all subsequent doses, to avoid intravascular or intrathecal injection. However, a negative aspiration for blood or cerebrospinal fluid does not ensure against an intravascular or intrathecal injection [see *Warnings and Precautions (5.9)*].
- Avoid rapid injection of a large volume of MARCAINE / MARCAINE WITH EPINEPHRINE and use fractional (incremental) doses when feasible.
- During major regional nerve blocks, such as those of the brachial plexus or lower extremity, the patient should have an indwelling intravenous catheter to assure adequate intravenous access. The lowest dosage of MARCAINE / MARCAINE WITH EPINEPHRINE that results in effective anesthesia should be used to avoid high

- plasma levels and serious adverse reactions.
- Perform careful and constant monitoring of cardiovascular and respiratory (adequacy of oxygenation and ventilation) vital signs and the patient's level of consciousness after each local anesthetic injection.
- Use MARCAINE WITH EPINEPHRINE in carefully restricted quantities in areas of the body supplied by end arteries or having otherwise compromised blood supply such as digits, nose, external ear, or penis [see *Warnings and Precautions (5.12)*].

2.2 Recommended Concentrations and Dosages of MARCAINE / MARCAINE WITH EPINEPHRINE

The dosage of MARCAINE / MARCAINE WITH EPINEPHRINE administered varies with the anesthetic procedure, the area to be anesthetized, the vascularity of the tissues, the number of neuronal segments to be blocked, the depth of anesthesia and degree of muscle relaxation required, the duration of anesthesia desired, individual tolerance, and the physical condition of the patient. Administer the smallest dosage and concentration required to produce the desired result.

The types of block and recommended MARCAINE / MARCAINE WITH EPINEPHRINE concentrations are shown in Table 1.

Table 1. Types of Block and Recommended MARCAINE / MARCAINE WITH EPINEPHRINE Concentrations

Type of Block	MARCAINE			MARCAINE WITH EPINEPHRINE	
	0.25% (2.5 mg/mL)	0.5% (5 mg/mL)	0.75% (7.5 mg/mL)*	0.25% (2.5 mg/mL)	0.5% (5 mg/mL)
Local infiltration	✓			✓	
Peripheral nerve block	✓	✓		✓	✓
Retrobulbar block			✓		
Sympathetic block	✓				
Caudal block†	✓	✓		✓	✓
Lumbar epidural block†	✓	✓	✓ (not for obstetrical anesthesia)	✓	✓
Epidural test dose					✓
Dental block					✓

✓ = indicated use [see *Warnings and Precautions (5.1)*].

* MARCAINE 0.75% (7.5 mg/mL) is not recommended for nonobstetrical surgical procedures in pregnant patients.

† Avoid use of multiple-dose vials of MARCAINE and MARCAINE WITH EPINEPHRINE for caudal or epidural anesthesia [see *Warnings and Precautions (5.4)*].

At recommended dosages, MARCAINE / MARCAINE WITH EPINEPHRINE produces complete sensory block, but the effect on motor function differs among the three concentrations. Table 2 provides information on the expected effect on motor function

for the three concentrations.

Table 2. MARCAINE / MARCAINE WITH EPINEPHRINE Concentration vs. Motor Function

MARCAINE Concentration	Motor Function
0.25% (2.5 mg/mL)*	When used for caudal, epidural, or peripheral nerve block, produces incomplete motor block. Should be used for operations in which muscle relaxation is not important, or when another means of providing muscle relaxation is used concurrently. Onset of action may be slower than with the 0.5% (5 mg/mL) or 0.75% (7.5 mg/mL) solutions.
0.5% (5 mg/mL)*	Provides motor blockade for caudal, epidural, or nerve block, but muscle relaxation may be inadequate for operations in which complete muscle relaxation is essential.
0.75% (7.5 mg/mL)†	Produces complete motor block. Most useful for epidural block in abdominal operations requiring complete muscle relaxation, and for retrobulbar anesthesia. Not for obstetrical anesthesia.

* These products include MARCAINE and MARCAINE WITH EPINEPHRINE [the epinephrine concentration (1:200,000) is not included in the table].

† These are only MARCAINE products [there is no 0.75% (7.5 mg/mL) concentration for MARCAINE WITH EPINEPHRINE].

The duration of anesthesia with MARCAINE / MARCAINE WITH EPINEPHRINE is such that for most indications, a single dose is sufficient.

The maximum dosage limit within the recommended dosage range must be individualized in each case after evaluating the size and physical status of the patient, as well as the anticipated rate of systemic absorption from a particular injection site.

The dosages in Table 3 are recommended as a guide for use in the average adult. These doses may be repeated once every three hours. Do not exceed a total daily dosage of 400 mg in 24 hours. The duration of anesthetic effect may be prolonged by the addition of epinephrine.

Table 3. Recommended Concentrations and Doses of MARCAINE / MARCAINE WITH EPINEPHRINE in Adults

Type of Block	Concentration of MARCAINE	Each Dose		Motor Block*
		mL	mg of MARCAINE	
Local infiltration	0.25% (2.5 mg/mL)†	Up to 70 (without epinephrine)	Up to 175 (without epinephrine)	—
		Up to 90 (with epinephrine)	Up to 225 (with epinephrine)	
	0.5% (5 mg/mL)†	5-35 (without epinephrine)	25-175 (without epinephrine)	moderate to complete

Peripheral nerve block		5-45 (with epinephrine)	25-225 (with epinephrine)	
	0.25% (2.5 mg/mL) [†]	5-70 (without epinephrine)	12.5-175 (without epinephrine)	moderate to complete
		5-90 (with epinephrine)	12.5-225 (with epinephrine)	
Retrobulbar block <i>[see Dosage and Administration (2.6)]</i>	0.75% (7.5 mg/mL)	2-4	15-30	complete
Sympathetic block	0.25% (2.5 mg/mL)	20-50	50-125	—
Caudal block <i>[see Dosage and Administration (2.4)]</i>	0.5% (5 mg/mL) [†]	15-30	75-150	moderate to complete
	0.25% (2.5 mg/mL) [†]	15-30	37.5-75	moderate
Lumbar epidural block <i>[see Dosage and Administration (2.3)]</i>	0.75% (7.5 mg/mL) [‡]	10-20	75-150	complete
	0.5% (5 mg/mL) [†]	10-20	50-100	moderate to complete
	0.25% (2.5 mg/mL) [†]	10-20	25-50	partial to moderate
Epidural test dose <i>[see Dosage and Administration (2.4)]</i>	0.5% (5 mg/mL) with epinephrine	2-3	10-15 (10-15 micrograms epinephrine)	—
Dental <i>[see Dosage and Administration (2.5)]</i>	0.5% (5 mg/mL) with epinephrine	1.8-3.6 per site	9-18 per site	—

* With continuous (intermittent) techniques, repeat doses increase the degree of motor block. The first repeat dose of 0.5% (5 mg/mL) may produce complete motor block. Intercostal nerve block with 0.25% (2.5 mg/mL) also may produce complete motor block for intra-thoracic and upper intra-abdominal surgery.

† Solutions with or without epinephrine (i.e., applies to MARCAINE and MARCAINE WITH EPINEPHRINE). The MARCAINE WITH EPINEPHRINE products include epinephrine (1:200,000).

‡ For single-dose use; not for intermittent epidural technique. Not for obstetrical anesthesia.

2.3 Use in Epidural Anesthesia

During the administration of epidural anesthesia, it is recommended that a test dose of

MARCAINE WITH EPINEPHRINE without antimicrobial preservative (0.5% bupivacaine with 1:200,000 epinephrine) be administered initially and the effects monitored before the full dose is given. When using a "continuous" catheter technique, test doses should be given prior to both the initial and all supplemental doses, because a catheter in the epidural space can migrate into a blood vessel or through the dura [see *Dosage and Administration (2.4)*].

During epidural administration, administer MARCAINE / MARCAINE WITH EPINEPHRINE, 0.5% (5 mg/mL) and MARCAINE 0.75% (7.5 mg/mL) solutions in incremental doses of 3 mL to 5 mL with sufficient time between doses to detect toxic manifestations of unintentional intravascular or intrathecal injection. Administer injections slowly, with frequent aspirations before and during the injection to avoid intravascular injection. Perform syringe aspirations before and during each supplemental injection in continuous (intermittent) catheter techniques. In obstetrics, use ONLY the 0.5% (5 mg/mL) and 0.25% (2.5 mg/mL) concentrations of MARCAINE / MARCAINE WITH EPINEPHRINE [see *Warnings and Precautions (5.1)*]; incremental doses of 3 mL to 5 mL of the 0.5% (5 mg/mL) solution not exceeding 50 mg to 100 mg at any dosing interval are recommended. Repeat doses should be preceded by a test dose containing epinephrine if not clinically contraindicated. Use only the single-dose vials for caudal or epidural anesthesia; avoid use of the multiple-dose vials for these procedures, which contain a preservative [see *Dosage and Administration (2.1, 2.4)*, *Warnings and Precautions (5.4, 5.9)*].

2.4 Test Dose for Caudal and Lumbar Epidural Blocks

Three mL of MARCAINE WITH EPINEPHRINE without antimicrobial preservative (0.5% bupivacaine with 1:200,000 epinephrine) is recommended for use as a test dose prior to caudal and lumbar epidural blocks when clinical conditions permit. This test dose may serve as a warning of unintended intravascular or intrathecal injection. Closely monitor for early clinical signs of toxicity following each test dose [see *Warnings and Precautions (5.9)*]. Allot adequate time for onset of spinal block to detect possible intrathecal injection. An intravascular or intrathecal injection is still possible even if results of the test dose are negative. The test dose itself may produce a systemic toxic reaction, high spinal, or cardiovascular effects from the epinephrine [see *Warnings and Precautions (5.2, 5.9)*, *Overdosage (10)*].

2.5 Use in Dentistry

MARCAINE WITH EPINEPHRINE 0.5% (5 mg/mL) is recommended for infiltration and block injection in the maxillary and mandibular area when a longer duration of local anesthesia is desired, such as for procedures generally associated with significant postoperative pain. The average dose of 1.8 mL (9 mg) per injection site will usually suffice; an occasional second dose of 1.8 mL (9 mg) may be used if necessary to produce adequate anesthesia after allowing 2 to 10 minutes for block onset [see *Clinical Pharmacology (12.2)*]. Use the lowest effective dose and allow time between injections; it is recommended that the total dose for all injection sites, spread out over a single dental sitting, not exceed 90 mg for a healthy adult patient (ten 1.8 mL injections of 0.5% (5 mg/mL) MARCAINE WITH EPINEPHRINE). Inject slowly and with frequent aspirations.

2.6 Use in Ophthalmic Surgery

When MARCAINE 0.75% (7.5 mg/mL) is used for retrobulbar block, complete corneal

anesthesia usually precedes onset of clinically acceptable external ocular muscle akinesia. Therefore, presence of akinesia rather than anesthesia alone should determine readiness of the patient for surgery [see *Warnings and Precautions (5.15)*].

3 DOSAGE FORMS AND STRENGTHS

MARCAINE (bupivacaine hydrochloride) injection is a clear, colorless solution available as:

- 0.25% (25 mg/10 mL) (2.5 mg/mL) in single-dose vials.
- 0.25% (75 mg/30 mL) (2.5 mg/mL) in single-dose vials.
- 0.25% (125 mg/50 mL) (2.5 mg/mL) in multiple-dose vials.
- 0.5% (50 mg/10 mL) (5 mg/mL) in single-dose vials.
- 0.5% (150 mg/30 mL) (5 mg/mL) in single-dose vials.
- 0.5% (250 mg/50 mL) (5 mg/mL) in multiple-dose vials.
- 0.75% (75 mg/10 mL) (7.5 mg/mL) in single-dose vials.
- 0.75% (225 mg/30 mL) (7.5 mg/mL) in single-dose vials.

MARCAINE with 1:200,000 epinephrine (bupivacaine hydrochloride and epinephrine) injection is a clear, colorless solution available as:

- 0.25% (25 mg/10 mL) (2.5 mg/mL) in single-dose vials.
- 0.25% (75 mg/30 mL) (2.5 mg/mL) in single-dose vials.
- 0.25% (125 mg/50 mL) (2.5 mg/mL) in multiple-dose vials.
- 0.5% (50 mg/10 mL) (5 mg/mL) in single-dose vials.
- 0.5% (150 mg/30 mL) (5 mg/mL) in single-dose vials.
- 0.5% (250 mg/50 mL) (5 mg/mL) in multiple-dose vials.

4 CONTRAINDICATIONS

MARCAINE / MARCAINE WITH EPINEPHRINE is contraindicated in:

- obstetrical paracervical block anesthesia. Its use in this technique has resulted in fetal bradycardia and death.
- intravenous regional anesthesia (Bier Block) [see *Warnings and Precautions (5.7)*].
- patients with a known hypersensitivity to bupivacaine or to any local anesthetic agent of the amide-type or to other components of MARCAINE / MARCAINE WITH EPINEPHRINE.

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Cardiac Arrest with Use of MARCAINE in Obstetrical Anesthesia

There have been reports of cardiac arrest with difficult resuscitation or death during use of MARCAINE for epidural anesthesia in obstetrical patients. In most cases, this has followed use of the 0.75% (7.5 mg/mL) concentration. Resuscitation has been difficult or impossible despite apparently adequate preparation and appropriate management. Cardiac arrest has occurred after convulsions resulting from systemic toxicity, presumably following unintentional intravascular injection. The 0.75% (7.5 mg/mL) concentration of MARCAINE is not recommended for obstetrical anesthesia and should

be reserved for surgical procedures where a high degree of muscle relaxation and prolonged effect are necessary.

5.2 Dose-Related Toxicity

The safety and effectiveness of MARCAINE / MARCAINE WITH EPINEPHRINE depend on proper dosage, correct technique, adequate precautions, and readiness for emergencies. Careful and constant monitoring of cardiovascular and respiratory (adequacy of ventilation) vital signs and the patient's state of consciousness should be performed after injection of MARCAINE / MARCAINE WITH EPINEPHRINE solutions.

Possible early warning signs of central nervous system (CNS) toxicity are restlessness, anxiety, incoherent speech, lightheadedness, numbness and tingling of the mouth and lips, metallic taste, tinnitus, dizziness, blurred vision, tremors, twitching, CNS depression, or drowsiness. Delay in proper management of dose-related toxicity, underventilation from any cause, and/or altered sensitivity may lead to the development of acidosis, cardiac arrest, and, possibly, death.

During major regional nerve blocks, such as those of the brachial plexus or lower extremity, the patient should have an indwelling intravenous catheter to assure adequate intravenous access. Use the lowest dosage of MARCAINE / MARCAINE WITH EPINEPHRINE that results in effective anesthesia to avoid high plasma levels and serious adverse effects. Avoid rapid injection of a large volume of MARCAINE / MARCAINE WITH EPINEPHRINE solution and administer fractional (incremental) doses when feasible.

Injection of repeated doses of MARCAINE / MARCAINE WITH EPINEPHRINE may cause significant increases in plasma levels with each repeated dose due to slow accumulation of the drug or its metabolites, or to slow metabolic degradation. Tolerance to elevated blood levels varies with the status of the patient. Debilitated, elderly patients and acutely ill patients should be given reduced doses commensurate with their age and physical status.

5.3 Methemoglobinemia

Cases of methemoglobinemia have been reported in association with local anesthetic use. Although all patients are at risk for methemoglobinemia, patients with glucose-6-phosphate dehydrogenase deficiency, congenital or idiopathic methemoglobinemia, cardiac or pulmonary compromise, infants under 6 months of age, and concurrent exposure to oxidizing agents or their metabolites are more susceptible to developing clinical manifestations of the condition [see *Drug Interactions (7.5)*]. If local anesthetics must be used in these patients, close monitoring for symptoms and signs of methemoglobinemia is recommended.

Signs of methemoglobinemia may occur immediately or may be delayed some hours after exposure, and are characterized by a cyanotic skin discoloration and/or abnormal coloration of the blood. Methemoglobin levels may continue to rise; therefore, immediate treatment is required to avert more serious CNS and cardiovascular adverse effects, including seizures, coma, arrhythmias, and death. Discontinue MARCAINE / MARCAINE WITH EPINEPHRINE and any other oxidizing agents. Depending on the severity of the signs and symptoms, patients may respond to supportive care, i.e., oxygen therapy, hydration. A more severe clinical presentation may require treatment with methylene blue, exchange transfusion, or hyperbaric oxygen.

5.4 Antimicrobial Preservatives in Multiple-Dose Vials

Avoid use of MARCAINE / MARCAINE WITH EPINEPHRINE solutions containing antimicrobial preservatives, i.e., those supplied in multiple-dose vials, for epidural or caudal anesthesia because safety has not been established with such use.

5.5 Chondrolysis with Intra-Articular Infusion

Intra-articular infusions of local anesthetics including MARCAINE following arthroscopic and other surgical procedures is an unapproved use, and there have been post-marketing reports of chondrolysis in patients receiving such infusions. The majority of reported cases of chondrolysis have involved the shoulder joint; cases of gleno-humeral chondrolysis have been described in pediatric and adult patients following intra-articular infusions of local anesthetics with and without epinephrine for periods of 48 to 72 hours. There is insufficient information to determine whether shorter infusion periods are associated with chondrolysis. The time of onset of symptoms, such as joint pain, stiffness and loss of motion can be variable, but may begin as early as the 2nd month after surgery. Currently, there is no effective treatment for chondrolysis; patients who experienced chondrolysis have required additional diagnostic and therapeutic procedures and some required arthroplasty or shoulder replacement.

5.6 Risk of Adverse Reactions Due to Drug Interactions with MARCAINE WITH EPINEPHRINE

Risk of Severe, Persistent Hypertension Due to Drug Interactions Between MARCAINE WITH EPINEPHRINE and Monoamine Oxidase Inhibitors and Tricyclic Antidepressants

Administration of MARCAINE WITH EPINEPHRINE (containing a vasoconstrictor) in patients receiving monoamine oxidase inhibitors (MAOI) or tricyclic antidepressants may result in severe, prolonged hypertension. Concurrent use of these agents should generally be avoided. In situations when concurrent therapy is necessary, careful monitoring of the patient's hemodynamic status is essential [*see Drug Interactions (7.2)*].

Risk of Severe, Persistent Hypertension or Cerebrovascular Accidents Due to Drug Interactions Between MARCAINE WITH EPINEPHRINE and Ergot-Type Oxytocic Drugs

Concurrent administration of MARCAINE WITH EPINEPHRINE and ergot-type oxytocic drugs may cause severe, persistent hypertension or cerebrovascular accidents. Avoid use of MARCAINE WITH EPINEPHRINE concomitantly with ergot-type oxytocic drugs [*see Drug Interactions (7.3)*].

Risk of Hypertension and Bradycardia Due to Drug Interactions Between MARCAINE WITH EPINEPHRINE and Nonselective Beta-Adrenergic Antagonists

Administration of MARCAINE WITH EPINEPHRINE (containing a vasoconstrictor) in patients receiving nonselective beta-adrenergic antagonists may cause severe hypertension and bradycardia. Concurrent use of these agents should generally be avoided. In situations when concurrent therapy is necessary, careful monitoring of the patient's blood pressure and heart rate is essential [*see Drug Interactions (7.4)*].

5.7 Risk of Cardiac Arrest with Intravenous Regional Anesthesia Use (Bier Block)

There have been reports of cardiac arrest and death during the use of bupivacaine for intravenous regional anesthesia (Bier Block). Information on safe dosages and techniques of administration of MARCAINE in this procedure is lacking. Therefore, MARCAINE / MARCAINE WITH EPINEPHRINE is contraindicated for use with this technique [see *Contraindications (4)*].

5.8 Allergic-Type Reactions to Sulfites in MARCAINE WITH EPINEPHRINE

MARCAINE WITH EPINEPHRINE contains sodium metabisulfite, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people. MARCAINE without epinephrine does not contain sodium metabisulfite.

5.9 Risk of Systemic Toxicities with Unintended Intravascular or Intrathecal Injection

Unintended intravascular or intrathecal injection of MARCAINE / MARCAINE WITH EPINEPHRINE may be associated with systemic toxicities, including CNS or cardiorespiratory depression and coma, progressing ultimately to respiratory arrest. Unintentional intrathecal injection during the intended performance of caudal or lumbar epidural block or nerve blocks near the vertebral column has resulted in underventilation or apnea ("Total or High Spinal"). A high spinal has been characterized by paralysis of the legs, loss of consciousness, respiratory paralysis, and bradycardia [see *Adverse Reactions (6)*].

Aspirate for blood or cerebrospinal fluid (where applicable) before injecting MARCAINE / MARCAINE WITH EPINEPHRINE, both the initial dose and all subsequent doses, to avoid intravascular or intrathecal injection. However, a negative aspiration for blood or cerebrospinal fluid does not ensure against an intravascular or intrathecal injection.

Use of Test Dose with Epidural Anesthesia

To serve as a warning of unintended intravascular or intrathecal injection, 3 mL of MARCAINE WITH EPINEPHRINE without antimicrobial preservative (0.5% bupivacaine with 1:200,000 epinephrine) may be used as a test dose prior to administration of the full dose in caudal and lumbar epidural blocks [see *Dosage and Administration (2.4)*]. Three mL of MARCAINE WITH EPINEPHRINE without antimicrobial preservative (0.5% bupivacaine with 1:200,000 epinephrine) contains 15 mg bupivacaine and 15 mcg epinephrine. An intravascular or intrathecal injection is still possible even if results of the test dose are negative.

Signs/symptoms of unintended intravascular or intrathecal injection of the test dose of MARCAINE WITH EPINEPHRINE and monitoring recommendations are described below.

- Unintended *intravascular* injection: Likely to produce a transient "epinephrine response" within 45 seconds, consisting of an increase in heart rate and/or systolic blood pressure, circumoral pallor, palpitations, and nervousness in the unsedated patient. The sedated patient may exhibit only a pulse rate increase of 20 or more beats per minute for 15 or more seconds. Therefore, following the test dose, the heart rate should be monitored for increases. Patients on beta-blockers may not manifest changes in heart rate, but blood pressure monitoring can detect a

transient rise in systolic blood pressure.

- Unintended *intrathecal* injection: Evidenced within a few minutes by signs of spinal block (e.g., decreased sensation of the buttocks, paresis of the legs, or, in the sedated patient, absent knee jerk).

The test dose itself may produce a systemic toxic reaction, high spinal or epinephrine-induced cardiovascular effects [see *Overdosage (10)*].

5.10 Risk of Toxicity in Patients with Hepatic Impairment

Because amide local anesthetics such as bupivacaine are metabolized by the liver, consider reduced dosing and increased monitoring for bupivacaine systemic toxicity in patients with moderate to severe hepatic impairment who are treated with MARCAINE / MARCAINE WITH EPINEPHRINE, especially with repeat doses [see *Use in Specific Populations (8.6)*].

5.11 Risk of Use in Patients with Impaired Cardiovascular Function

MARCAINE / MARCAINE WITH EPINEPHRINE should be given in reduced doses in patients with impaired cardiovascular function (e.g., hypotension, heartblock) because they may be less able to compensate for functional changes associated with the prolongation of AV conduction produced by MARCAINE / MARCAINE WITH EPINEPHRINE. Monitor patients closely for blood pressure, heart rate, and ECG changes.

5.12 Risk of Ischemic Injury or Necrosis in Body Areas with Limited Blood Supply

Use MARCAINE WITH EPINEPHRINE in carefully restricted quantities in areas of the body supplied by end arteries or having otherwise compromised blood supply such as digits, nose, external ear, or penis. Patients with hypertensive vascular disease may exhibit exaggerated vasoconstrictor response. Ischemic injury or necrosis may result.

5.13 Risk of Cardiac Arrhythmias with Concomitant Use of Potent Inhalation Anesthetics

Serious dose-related cardiac arrhythmias may occur if preparations containing a vasoconstrictor such as epinephrine (e.g., MARCAINE WITH EPINEPHRINE) are used in patients during or following the administration of potent inhalation anesthetics [see *Drug Interactions (7.6)*]. In deciding whether to concurrently use MARCAINE WITH EPINEPHRINE with potent inhalation anesthetics in the same patient, the combined action of both agents upon the myocardium, the concentration and volume of vasoconstrictor used, and the time since injection, when applicable, should be taken into account.

5.14 Risk of Adverse Reactions with Use in Head and Neck Area

Small doses of local anesthetics (e.g., MARCAINE) injected into the head and neck area, including retrobulbar, dental, and stellate ganglion blocks, may produce adverse reactions similar to systemic toxicity seen with unintentional intravascular injections of larger doses. The injection procedures require the utmost care. Confusion, convulsions, respiratory depression, and/or respiratory arrest, and cardiovascular stimulation or depression have been reported. These reactions may be due to intra-arterial injection of the local anesthetic with retrograde flow to the cerebral circulation. They may also be

due to puncture of the dural sheath of the optic nerve during retrobulbar block with diffusion of any local anesthetic along the subdural space to the midbrain. Monitor circulation and respiration and constantly observe patients receiving MARCAINE / MARCAINE WITH EPINEPHRINE blocks. Resuscitative equipment and drugs, and personnel for treating adverse reactions should be immediately available. Dosage recommendations should not be exceeded [see *Dosage and Administration (2.2)*].

5.15 Risk of Respiratory Arrest with Use in Ophthalmic Surgery

Clinicians who perform retrobulbar blocks should be aware that there have been reports of respiratory arrest following local anesthetic injection. Prior to retrobulbar block (e.g., with MARCAINE / MARCAINE WITH EPINEPHRINE), as with all other regional procedures, resuscitative equipment and drugs, and personnel to manage respiratory arrest or depression, convulsions, and cardiac stimulation or depression should be immediately available [see *Warnings and Precautions (5.14)*]. As with other anesthetic procedures, patients should be constantly monitored following ophthalmic blocks for signs of these adverse reactions, which may occur following relatively low total doses.

A concentration of 0.75% bupivacaine is indicated for retrobulbar block; however, this concentration is not indicated for any other peripheral nerve block, including the facial nerve, and not indicated for local infiltration, including the conjunctiva [see *Indications and Usage (1)*].

5.16 Risk of Inadvertent Trauma to Tongue, Lips, and Buccal Mucosa in Dental Applications

Because of the long duration of anesthesia, when MARCAINE WITH EPINEPHRINE [0.5% (5 mg/mL) of bupivacaine] is used for dental injections, warn patients about the possibility of inadvertent trauma to tongue, lips, and buccal mucosa and advise them not to chew solid foods until sensation returns [see *Patient Counseling Information (17)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions have been reported and described in the Warnings and Precautions section of the labeling:

- Cardiac Arrest in Obstetrical Anesthesia [see *Warnings and Precautions (5.1)*]
- Dose-Related Toxicity [see *Warnings and Precautions (5.2)*]
- Methemoglobinemia [see *Warnings and Precautions (5.3)*]
- Chondrolysis with Intra-Articular Infusion [see *Warnings and Precautions (5.5)*]
- Severe, Persistent Hypertension, Cerebrovascular Accidents, and Bradycardia Due to Drug Interactions [see *Warnings and Precautions (5.6)*]
- Cardiac Arrest with Intravenous Regional Anesthesia Use [see *Contraindications (4), Warnings and Precautions (5.7)*]
- Allergic-Type Reactions [see *Warnings and Precautions (5.8)*]
- Systemic Toxicities with Unintended Intravascular or Intrathecal Injection [see *Warnings and Precautions (5.9)*]
- Respiratory Arrest Following Retrobulbar Block [see *Warnings and Precautions (5.15)*]

The following adverse reactions from voluntary reports or clinical studies have been

reported with bupivacaine or bupivacaine and epinephrine. Because many of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Adverse reactions to MARCAINE / MARCAINE WITH EPINEPHRINE are characteristic of those associated with other amide-type local anesthetics. A major cause of adverse reactions to this group of drugs is excessive plasma levels, which may be due to overdosage, unintentional intravascular injection, or slow metabolic degradation.

The most commonly encountered acute adverse reactions that demand immediate counter-measures were related to the CNS and the cardiovascular system. These adverse reactions were generally dose-related and due to high plasma levels which may have resulted from overdosage, rapid absorption from the injection site, diminished tolerance, or from unintentional intravascular injection of the local anesthetic solution. In addition to systemic dose-related toxicity, unintentional intrathecal injection of drug during the intended performance of caudal or lumbar epidural block or nerve blocks near the vertebral column (especially in the head and neck region) has resulted in underventilation or apnea ("Total or High Spinal"). Also, hypotension due to loss of sympathetic tone and respiratory paralysis or underventilation due to cephalad extension of the motor level of anesthesia have occurred. This has led to secondary cardiac arrest when untreated.

Nervous System Disorders: Adverse reactions were characterized by excitation and/or depression of the central nervous system and included restlessness, anxiety, dizziness, tinnitus, blurred vision, tremors, convulsions, drowsiness, unconsciousness, respiratory arrest, nausea, vomiting, chills, pupillary constriction.

In the practice of caudal or lumbar epidural block, unintentional penetration of the subarachnoid space by the catheter or needle has occurred. Subsequent adverse effects may have depended partially on the amount of drug administered intrathecally and the physiological and physical effects of a dural puncture. A high spinal has been characterized by paralysis of the legs, loss of consciousness, respiratory paralysis, and bradycardia.

Neurologic effects following epidural or caudal anesthesia have included spinal block of varying magnitude (including high or total spinal block); hypotension secondary to spinal block; urinary retention; fecal and urinary incontinence; loss of perineal sensation and sexual function; persistent anesthesia, paresthesia, weakness, paralysis of the lower extremities and loss of sphincter control, all of which had slow, incomplete, or no recovery; headache; backache; septic meningitis; meningismus; slowing of labor; increased incidence of forceps delivery; and cranial nerve palsies due to traction on nerves from loss of cerebrospinal fluid.

Neurologic effects following other procedures or routes of administration have included persistent anesthesia, paresthesia, weakness, paralysis, all with slow, incomplete, or no recovery.

Convulsions: Incidence varied with the procedure used and the total dose administered. In a survey of studies of epidural anesthesia, overt toxicity progressing to convulsions occurred in approximately 0.1% of local anesthetic administrations. The incidences of adverse neurologic reactions associated with the use of local anesthetics may be related to the total dose of local anesthetic administered and are also dependent upon the

particular drug used, the route of administration, and the physical status of the patient.

Cardiac Disorders: High doses or unintentional intravascular injection have led to high plasma levels and related depression of the myocardium, decreased cardiac output, heartblock, hypotension, bradycardia, ventricular arrhythmias, including ventricular tachycardia and ventricular fibrillation, and cardiac arrest [see *Warnings and Precautions (5.9)*].

Immune System Disorders: Allergic-type reactions have occurred as a result of sensitivity to bupivacaine or to other formulation ingredients, such as the antimicrobial preservative methylparaben contained in multiple-dose vials or sulfites in epinephrine-containing solutions. These reactions were characterized by signs such as urticaria, pruritus, erythema, angioneurotic edema (including laryngeal edema), tachycardia, sneezing, nausea, vomiting, dizziness, syncope, excessive sweating, elevated temperature, and severe hypotension. Cross sensitivity among members of the amide-type local anesthetic group has been reported [see *Warnings and Precautions (5.8)*].

7 DRUG INTERACTIONS

7.1 Local Anesthetics

The toxic effects of local anesthetics are additive. If coadministration of other local anesthetics with MARCAINE / MARCAINE WITH EPINEPHRINE cannot be avoided, monitor patients for neurologic and cardiovascular effects related to local anesthetic systemic toxicity [see *Dosage and Administration (2.1)*, *Warnings and Precautions (5.2)*].

7.2 Monoamine Oxidase Inhibitors and Tricyclic Antidepressants

The administration of MARCAINE WITH EPINEPHRINE to patients receiving monoamine oxidase inhibitors, or tricyclic antidepressants may produce severe, prolonged hypertension. Concurrent use of these agents should generally be avoided. In situations when concurrent therapy is necessary, careful monitoring of the patient's hemodynamic status is essential [see *Warnings and Precautions (5.6)*].

7.3 Ergot-Type Oxytocic Drugs

Concurrent administration of MARCAINE WITH EPINEPHRINE and ergot-type oxytocic drugs may cause severe, persistent hypertension or cerebrovascular accidents. Avoid use of MARCAINE WITH EPINEPHRINE concomitantly with ergot-type oxytocic drugs [see *Warnings and Precautions (5.6)*].

7.4 Nonselective Beta-Adrenergic Antagonists

Administration of MARCAINE WITH EPINEPHRINE (containing a vasoconstrictor) in patients receiving nonselective beta-adrenergic antagonists may cause severe hypertension and bradycardia. Concurrent use of these agents should generally be avoided. In situations when concurrent therapy is necessary, careful monitoring of the patient's blood pressure and heart rate is essential [see *Warnings and Precautions (5.6)*].

7.5 Drugs Associated with Methemoglobinemia

Patients who are administered MARCAINE / MARCAINE WITH EPINEPHRINE are at

increased risk of developing methemoglobinemia when concurrently exposed to the following drugs, which could include other local anesthetics [see *Warnings and Precautions* (5.3)].

Examples of Drugs Associated with Methemoglobinemia:

Class	Examples
Nitrates/Nitrites	nitric oxide, nitroglycerin, nitroprusside, nitrous oxide
Local anesthetics	articaine, benzocaine, bupivacaine, lidocaine, mepivacaine, prilocaine, procaine, ropivacaine, tetracaine
Antineoplastic agents	cyclophosphamide, flutamide, hydroxyurea, ifosfamide, rasburicase
Antibiotics	dapsone, nitrofurantoin, para-aminosalicylic acid, sulfonamides
Antimalarials	chloroquine, primaquine
Anticonvulsants	phenobarbital, phenytoin, sodium valproate
Other drugs	acetaminophen, metoclopramide, quinine, sulfasalazine

7.6 Potent Inhalation Anesthetics

Serious dose-related cardiac arrhythmias may occur if preparations containing a vasoconstrictor such as epinephrine (e.g., MARCAINE WITH EPINEPHRINE) are used in patients during or following the administration of potent inhalation anesthetics [see *Warnings and Precautions* (5.13)].

7.7 Phenothiazines and Butyrophenones

Phenothiazines and butyrophenones may reduce or reverse the pressor effect of epinephrine. Concurrent use of MARCAINE WITH EPINEPHRINE and these agents should generally be avoided. In situations when concurrent therapy is necessary, careful patient monitoring is essential.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

MARCAINE / MARCAINE WITH EPINEPHRINE is contraindicated for obstetrical paracervical block anesthesia. Its use in this technique has resulted in fetal bradycardia and death [see *Contraindications* (4), *Warnings and Precautions* (5.1)].

There are no available data on use of MARCAINE / MARCAINE WITH EPINEPHRINE in pregnant women to inform a drug-associated risk of adverse developmental outcomes.

In animal studies, embryo-fetal lethality was noted when bupivacaine was administered subcutaneously to pregnant rabbits during organogenesis at clinically relevant doses. Decreased pup survival was observed in a rat pre- and post-natal developmental study (dosing from implantation through weaning) at a dose level comparable to the daily maximum recommended human dose (MRHD) on a body surface area (BSA) basis.

Based on animal data, advise pregnant women of the potential risks to a fetus (see *Data*).

Local anesthetics rapidly cross the placenta, and when used for epidural, caudal, or pudendal block anesthesia, can cause varying degrees of maternal, fetal, and neonatal toxicity [see *Clinical Pharmacology (12.3)*]. The incidence and degree of toxicity depend upon the procedure performed, the type, and amount of drug used, and the technique of drug administration. Adverse reactions in the parturient, fetus, and neonate involve alterations of the CNS, peripheral vascular tone, and cardiac function.

If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, inform the patient of the potential hazard to the fetus. The estimated background risk of major birth defects and miscarriage for the indicated populations are unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

Clinical Considerations

Maternal Adverse Reactions

Maternal hypotension has resulted from regional anesthesia. Local anesthetics produce vasodilation by blocking sympathetic nerves. The supine position is dangerous in pregnant women at term because of aortocaval compression by the gravid uterus. Therefore, during treatment of systemic toxicity, maternal hypotension or fetal bradycardia following regional block, the parturient should be maintained in the left lateral decubitus position if possible, or manual displacement of the uterus off the great vessels be accomplished. Elevating the patient's legs will also help prevent decreases in blood pressure. The fetal heart rate also should be monitored continuously and electronic fetal monitoring is highly advisable.

Labor or Delivery

Epidural, caudal, or pudendal anesthesia may alter the forces of parturition through changes in uterine contractility or maternal expulsive efforts. Epidural anesthesia has been reported to prolong the second stage of labor by removing the parturient's reflex urge to bear down or by interfering with motor function. The use of obstetrical anesthesia may increase the need for forceps assistance.

The use of some local anesthetic drug products during labor and delivery may be followed by diminished muscle strength and tone for the first day or two of life. This has not been reported with bupivacaine.

It is extremely important to avoid aortocaval compression by the gravid uterus during administration of regional block to parturients. To do this, the patient must be maintained in the left lateral decubitus position or a blanket roll or sandbag may be placed beneath the right hip and gravid uterus displaced to the left.

Data

Animal Data

Bupivacaine hydrochloride produced developmental toxicity when administered subcutaneously to pregnant rats and rabbits at clinically relevant doses.

Bupivacaine hydrochloride was administered subcutaneously to rats at doses of 4.4,

13.3, & 40 mg/kg and to rabbits at doses of 1.3, 5.8, & 22.2 mg/kg during the period of organogenesis (implantation to closure of the hard palate). The high doses are comparable to the daily MRHD of 400 mg/day on a mg/m² BSA basis. No embryo-fetal effects were observed in rats at the high dose which caused increased maternal lethality. An increase in embryo-fetal deaths was observed in rabbits at the high dose in the absence of maternal toxicity with the fetal No Observed Adverse Effect Level representing approximately 0.3 times the MRHD on a BSA basis.

In a rat pre- and post-natal developmental study (dosing from implantation through weaning) conducted at subcutaneous doses of 4.4, 13.3, & 40 mg/kg, decreased pup survival was observed at the high dose. The high dose is comparable to the daily MRHD of 400 mg/day on a BSA basis.

8.2 Lactation

Risk Summary

Lactation studies have not been conducted with bupivacaine. Bupivacaine has been reported to be excreted in human milk suggesting that the nursing infant could be theoretically exposed to a dose of the drug. MARCAINE / MARCAINE WITH EPINEPHRINE should be administered to lactating women only if clearly indicated. Studies assessing the effects of MARCAINE / MARCAINE WITH EPINEPHRINE in breastfed children have not been performed. Studies to assess the effect of MARCAINE / MARCAINE WITH EPINEPHRINE on milk production or excretion have not been performed. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for bupivacaine and any potential adverse effects on the breastfed child from bupivacaine or from the underlying maternal condition.

8.4 Pediatric Use

MARCAINE / MARCAINE WITH EPINEPHRINE is approved for use in adults. Administration of MARCAINE / MARCAINE WITH EPINEPHRINE in pediatric patients younger than 12 years is not recommended.

Continuous infusions of bupivacaine in pediatric patients have been reported to result in high systemic levels of bupivacaine and seizures; high plasma levels may also be associated with cardiovascular abnormalities.

8.5 Geriatric Use

Patients 65 years and over, particularly those with hypertension, may be at increased risk for developing hypotension while undergoing anesthesia with MARCAINE / MARCAINE WITH EPINEPHRINE.

In clinical studies of bupivacaine, elderly patients reached the maximal spread of analgesia and maximal motor blockade more rapidly than younger adult patients.

Differences in various pharmacokinetic parameters have been observed between elderly and younger adult patients [*see Clinical Pharmacology (12.3)*].

This product is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. Elderly patients may

require lower doses of MARCAINE / MARCAINE WITH EPINEPHRINE.

8.6 Hepatic Impairment

Amide-type local anesthetics, such as bupivacaine, are metabolized by the liver. Patients with severe hepatic impairment, because of their inability to metabolize local anesthetics normally, are at a greater risk of developing toxic plasma concentrations, and potentially local anesthetic systemic toxicity. Therefore, consider reduced dosing and increased monitoring for local anesthetic systemic toxicity in patients with moderate to severe hepatic impairment treated with MARCAINE / MARCAINE WITH EPINEPHRINE, especially with repeat doses [see *Warnings and Precautions (5.10)*].

8.7 Renal Impairment

Bupivacaine is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with renal impairment. This should be considered when selecting the MARCAINE / MARCAINE WITH EPINEPHRINE dosage [see *Use in Specific Populations (8.5)*].

10 OVERDOSAGE

Clinical Presentation

Acute emergencies from use of MARCAINE / MARCAINE WITH EPINEPHRINE are generally related to high plasma levels encountered during therapeutic use or to unintended intrathecal injection [see *Warnings and Precautions (5.2, 5.9), Adverse Reactions (6)*].

If not treated immediately, convulsions with simultaneous hypoxia, hypercarbia, and acidosis plus myocardial depression from the direct effects of bupivacaine may result in cardiac arrhythmias, bradycardia, asystole, ventricular fibrillation, or cardiac arrest. Respiratory abnormalities, including apnea, may occur. Hypoventilation or apnea due to unintentional intrathecal injection of MARCAINE / MARCAINE WITH EPINEPHRINE may produce these same signs and also lead to cardiac arrest if ventilatory support is not instituted. If cardiac arrest should occur, successful outcome may require prolonged resuscitative efforts.

Management

The first step in the management of systemic toxic reactions, as well as hypoventilation or apnea due to unintentional intrathecal injection of MARCAINE / MARCAINE WITH EPINEPHRINE, consists of immediate attention to the establishment and maintenance of a patent airway and effective assisted or controlled ventilation with 100% oxygen with a delivery system capable of permitting immediate positive airway pressure by mask. Endotracheal intubation, using drugs and techniques familiar to the clinician, may be indicated after initial administration of oxygen by mask if difficulty is encountered in the maintenance of a patent airway, or if prolonged ventilatory support (assisted or controlled) is indicated.


If necessary, use drugs to manage the convulsions. A bolus intravenous dose of a benzodiazepine will counteract CNS stimulation related to MARCAINE. Immediately after the institution of ventilatory measures, evaluate the adequacy of the circulation. Supportive treatment of circulatory depression may require Advanced Cardiac Life

Support measures.

11 DESCRIPTION


MARCAINE / MARCAINE WITH EPINEPHRINE contains bupivacaine hydrochloride, an amide local anesthetic, as the active pharmaceutical ingredient. The route of administration for MARCAINE (without epinephrine) is by injection, for infiltration, perineural, caudal, epidural, or retrobulbar use. The route of administration for MARCAINE WITH EPINEPHRINE is by injection, for infiltration, perineural, caudal, or epidural use. Multiple-dose vials contain methylparaben [see *Warnings and Precautions* (5.4)].

Bupivacaine hydrochloride is 2-piperidinecarboxamide, 1-butyl-N-(2,6-dimethylphenyl)-, monohydrochloride, monohydrate. It is a white crystalline powder that is freely soluble in 95 percent ethanol, soluble in water, and slightly soluble in chloroform or acetone. It has the following structural formula:



MARCAINE with 1:200,000 epinephrine, contains bupivacaine hydrochloride and epinephrine bitartrate (an alpha and beta-adrenergic agonist) as active pharmaceutical ingredients. This product is for injection via local infiltration, peripheral nerve block, and caudal and lumbar epidural blocks. Multiple dose vials contain methylparaben and they should not be used for caudal and lumbar epidural blocks.

Epinephrine bitartrate is (-) - 3,4-Dihydroxy- α -[(methylamino)methyl] benzyl alcohol (+)-tartrate (1:1) salt. Epinephrine is a vasoconstrictor. It has the following structural formula:



MARCAINE (bupivacaine hydrochloride) is a clear and colorless sterile isotonic solution. Each mL of single-dose vial contains 2.5 mg, 5 mg or 7.5 mg of bupivacaine hydrochloride (equivalent to 2.22 mg, 4.44 mg or 6.66 mg of bupivacaine, respectively), sodium chloride for isotonicity, sodium hydroxide or hydrochloric acid to adjust the pH between 4 and 6.5, in water for injection.

For the Multiple-dose vials, each mL also contains 1 mg methylparaben as preservative.

MARCAINE with epinephrine 1:200,000 (as bitartrate) is a clear and colorless sterile isotonic solution. Each mL contains 2.5 mg or 5 mg bupivacaine hydrochloride (equivalent to 2.22 mg or 4.44 mg of bupivacaine, respectively), and 0.0091 mg epinephrine bitartrate (equivalent to 0.005 mg of epinephrine), with sodium chloride for isotonicity, 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, and 2 mg ascorbic acid as antioxidants, 0.0017 mL 60% sodium lactate buffer, and 0.1 mg edetate calcium disodium as stabilizer. The pH of these solutions is adjusted to between 3.4 and 4.5 with sodium hydroxide or hydrochloric acid.

For the Multiple-dose vials, each mL also contains 1 mg methylparaben as preservative.

The specific gravity of *MARCAINE 0.5% with epinephrine 1:200,000* (as bitartrate) at 25°C is 1.008 and at 37°C is 1.008.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Bupivacaine blocks the generation and the conduction of nerve impulses, presumably by increasing the threshold for electrical excitation in the nerve, by slowing the propagation of the nerve impulse, and by reducing the rate of rise of the action potential. In general, the progression of anesthesia is related to the diameter, myelination, and conduction velocity of affected nerve fibers. Clinically, the order of loss of nerve function is as follows: (1) pain, (2) temperature, (3) touch, (4) proprioception, and (5) skeletal muscle tone.

Epinephrine is a vasoconstrictor added to bupivacaine to slow absorption into the general circulation and thus prolong maintenance of an active tissue concentration.

12.2 Pharmacodynamics

Systemic absorption of bupivacaine produces effects on the cardiovascular system and CNS. At blood concentrations achieved with normal therapeutic doses, changes in cardiac conduction, excitability, refractoriness, contractility, and peripheral vascular resistance are minimal. However, toxic blood concentrations depress cardiac conduction and excitability, which may lead to atrioventricular block, ventricular arrhythmias, and cardiac arrest, sometimes resulting in fatalities. In addition, myocardial contractility is depressed and peripheral vasodilation occurs, leading to decreased cardiac output and arterial blood pressure. These cardiovascular changes are more likely to occur after unintended intravascular injection of bupivacaine [see *Warnings and Precautions (5.9)*].

Following systemic absorption, bupivacaine can produce CNS stimulation, CNS depression, or both. Apparent central stimulation is manifested as restlessness, tremors, and shivering, progressing to convulsions, followed by CNS depression and coma progressing ultimately to respiratory arrest. However, bupivacaine has a primary depressant effect on the medulla and on higher centers. The depressed stage may occur without a prior excited state.

The duration of local anesthesia after administration of *MARCAINE* is longer than that observed after administration of other commonly used short-acting local anesthetics. There appears to be a period of analgesia that persists after the resolution of the block

and return of sensation.

The onset of action following dental injections is usually 2 to 10 minutes and may last up to 7 hours. The duration of anesthetic effect is prolonged by the addition of epinephrine 1:200,000.

12.3 Pharmacokinetics

Systemic plasma levels of bupivacaine following administration of MARCAINE do not correlate with local efficacy.

Absorption

The rate of systemic absorption of bupivacaine is dependent upon the total dose and concentration of drug administered, the route of administration, the vascularity of the administration site, and the presence or absence of epinephrine in the anesthetic solution. A dilute concentration of epinephrine (1:200,000) usually reduces the rate of absorption and peak plasma concentration of bupivacaine, permitting the use of moderately larger total doses and sometimes prolonging the duration of action [see *Dosage and Administration (2)*].

After injection of MARCAINE for caudal, epidural, or peripheral nerve block, peak levels of bupivacaine in the blood are reached in 30 to 45 minutes, followed by a decline to insignificant levels during the next three to six hours.

Distribution

Bupivacaine appears to cross the placenta by passive diffusion. The rate and degree of diffusion is governed by (1) the degree of plasma protein binding, (2) the degree of ionization, and (3) the degree of lipid solubility. Fetal/ maternal ratios of bupivacaine appear to be inversely related to the degree of plasma protein binding, because only the free, unbound drug is available for placental transfer. Bupivacaine with a high protein binding capacity (95%) has a low fetal/maternal ratio (0.2 to 0.4). The extent of placental transfer is also determined by the degree of ionization and lipid solubility of the drug. Lipid soluble, nonionized drugs readily enter the fetal blood from the maternal circulation.

Depending upon the route of administration, bupivacaine is distributed to some extent to all body tissues, with high concentrations found in highly perfused organs such as the liver, lungs, heart, and brain.

Pharmacokinetic studies on the plasma profile of bupivacaine after direct intravenous injection (MARCAINE is not approved for intravenous use) suggest a three-compartment open model. The first compartment is represented by the rapid intravascular distribution of the drug. The second compartment represents the equilibration of the drug throughout the highly perfused organs such as the brain, myocardium, lungs, kidneys, and liver. The third compartment represents an equilibration of the drug with poorly perfused tissues, such as muscle and fat.

Elimination

The half-life of bupivacaine in adults is 2.7 hours.

Metabolism

Amide-type local anesthetics such as bupivacaine are metabolized primarily in the liver via conjugation with glucuronic acid. Pipecoloxylidene is the major metabolite of

bupivacaine. The elimination of drug from tissue distribution depends largely upon the availability of binding sites in the circulation to carry it to the liver where it is metabolized.

Excretion

The kidney is the main excretory organ for most local anesthetics and their metabolites. Urinary excretion is affected by urinary perfusion and factors affecting urinary pH. Only 6% of bupivacaine is excreted unchanged in the urine.

Specific Populations

Geriatric Patients

Elderly patients exhibited higher peak plasma concentrations than younger patients following administration of MARCAINE. The total plasma clearance was decreased in these patients [*see Use in Specific Populations (8.5)*].

Patients with Hepatic Impairment

Various pharmacokinetic parameters of the local anesthetics can be significantly altered by the presence of hepatic disease. Patients with hepatic disease, especially those with severe hepatic disease, may be more susceptible to the potential toxicities of the amide-type local anesthetics [*see Use in Specific Populations (8.6)*].

Patients with Renal Impairment

Various pharmacokinetic parameters of the local anesthetics can be significantly altered by the presence of renal disease, factors affecting urinary pH, and renal blood flow [*see Use in Specific Populations (8.5, 8.7)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies in animals to evaluate the carcinogenic potential of bupivacaine hydrochloride have not been conducted.

Mutagenesis

The mutagenic potential of bupivacaine hydrochloride has not been determined.

Impairment of Fertility

The effect of bupivacaine on fertility has not been determined.

16 HOW SUPPLIED/STORAGE AND HANDLING

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature.]

MARCAINE—Solutions of MARCAINE that do not contain epinephrine may be autoclaved. Autoclave at 15-pound pressure, 121°C (250°F) for 15 minutes. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate.

Unit of Sale	Concentration
NDC 0409-1559-10 Tray of 10 single-dose vials	0.25% 25 mg/10 mL (2.5 mg/mL)
NDC 0409-2510-25 Tray of 25 single-dose vials	0.25% 25 mg/10 mL (2.5 mg/mL)
NDC 0409-1559-30 Carton of 10 single-dose vials	0.25% 75 mg/30 mL (2.5 mg/mL)
NDC 0409-7535-25 Tray of 25 single-dose vials	0.25% 75 mg/30 mL (2.5 mg/mL)
NDC 0409-1587-50 Carton of 1 multiple-dose vial	0.25% 125 mg/50 mL (2.5 mg/mL)
NDC 0409-1250-25 Tray of 25 multiple-dose vials	0.25% 125 mg/50 mL (2.5 mg/mL)
NDC 0409-1560-10 Tray of 10 single-dose vials	0.5% 50 mg/10 mL (5 mg/mL)
NDC 0409-5010-25 Tray of 25 single-dose vials	0.5% 50 mg/10 mL (5 mg/mL)
NDC 0409-1560-29 Carton of 10 single-dose vials	0.5% 150 mg/30 mL (5 mg/mL)
NDC 0409-1530-25 Tray of 25 single-dose vials	0.5% 150 mg/30 mL (5 mg/mL)
NDC 0409-1610-50 Carton of 1 multiple-dose vial	0.5% 250 mg/50 mL (5 mg/mL)
NDC 0409-0525-25 Tray of 25 multiple-dose vials	0.5% 250 mg/50 mL (5 mg/mL)
NDC 0409-1582-10 Tray of 10 single-dose vials	0.75% 75 mg/10 mL (7.5 mg/mL)
NDC 0409-7510-25 Tray of 25 single-dose vials	0.75% 75 mg/10 mL (7.5 mg/mL)
NDC 0409-1582-29 Carton of 10 single-dose vials	0.75% 225 mg/30 mL (7.5 mg/mL)
NDC 0409-2253-25	0.75%

Tray of 25 single-dose vials	225 mg/30 mL (7.5 mg/mL)
------------------------------	------------------------------------

For single-dose vials: Discard unused portion.

MARCAINE with epinephrine 1:200,000 (as bitartrate)—Do not autoclave solutions of *MARCAINE* that contain epinephrine and protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate.

Unit of Sale	Concentration
NDC 0409-1746-10 Carton of 10 single-dose vials	0.25% 25 mg/10 mL (2.5 mg/mL)
NDC 0409-1746-30 Carton of 10 single-dose vials	0.25% 75 mg/30 mL (2.5 mg/mL)
NDC 0409-1752-50 Carton of 1 multiple-dose vial	0.25% 125 mg/50 mL (2.5 mg/mL)
NDC 0409-1749-10 Carton of 10 single-dose vials	0.5% 50 mg/10 mL (5 mg/mL)
NDC 0409-1749-29 Carton of 10 single-dose vials	0.5% 150 mg/30 mL (5 mg/mL)
NDC 0409-1755-50 Carton of 1 multiple-dose vial	0.5% 250 mg/50 mL (5 mg/mL)

For single-dose vials: Discard unused portion.

17 PATIENT COUNSELING INFORMATION

Allergic-Type Reactions

Assess if the patient has had allergic-type reactions to amide-type local anesthetics or to other formulation ingredients, such as the antimicrobial preservative methylparaben contained in multiple-dose vials or sulfites in epinephrine-containing solutions [see *Contraindications (4)*, *Warnings and Precautions (5.8)*, *Adverse Reactions (6)*].

Temporary Loss of Sensation and Motor Activity After Caudal or Epidural Anesthesia

When appropriate, patients should be informed in advance that they may experience temporary loss of sensation and motor activity, usually in the lower half of the body, following proper administration of caudal or epidural anesthesia.

Instructions After Dental Injection of MARCAINE

Advise patients receiving dental injections of *MARCAINE* not to chew solid foods or to

test the anesthetized area by biting or probing until anesthesia has worn off (up to 7 hours) [see *Warnings and Precautions (5.16)*].

Methemoglobinemia

Inform patients that use of local anesthetics may cause methemoglobinemia, a serious condition that must be treated promptly. Advise patients or caregivers to seek immediate medical attention if they or someone in their care experience the following signs or symptoms: pale, gray, or blue colored skin (cyanosis); headache; rapid heart rate; shortness of breath; lightheadedness; or fatigue [see *Warnings and Precautions (5.3)*].

This product's labeling may have been updated. For the most recent prescribing information, please visit www.pfizer.com.

For medical information about MARCAINE / MARCAINE WITH EPINEPHRINE, please visit www.pfizermedinfo.com or call 1-800-438-1985.

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

LAB-1178-6.0

PRINCIPAL DISPLAY PANEL - 25 mg/10 mL Vial Label - 1559

10 mL
Single-dose Vial
NDC 0409-1559-18
Rx only


Preservative-Free

Marcaine™ 0.25%
bupivacaine HCl injection, USP
25 mg/10 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA


NOT FOR INTRATHECAL ANESTHESIA.

Hospira



PRINCIPAL DISPLAY PANEL - 25 mg/10 mL Vial Tray - 1559

10 mL
 Preservative-Free
 10 Single-dose Vials
 Rx only
 NDC 0409-1559-10
 Contains 10 of NDC 0409-1559-18
 Marcaine™ 0.25%
 bupivacaine HCl injection, USP
 25 mg/10 mL (2.5 mg/mL)
 For INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
 NOT FOR INTRATHECAL ANESTHESIA
 Hospira



PRINCIPAL DISPLAY PANEL - 75 mg/30 mL Vial Label - 1559

30 mL
 SINGLE-DOSE VIAL- PRESERVATIVE FREE
 NDC 0409-1559-19
 Rx only
 Marcaine™ 0.25%
 bupivacaine HCl injection, USP

75 mg/30 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK,
CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

Hospira

FPO DataBar Limited 10 MIL

FPO 2D 15 MIL
PAA160399

30 mL
SINGLE-DOSE VIAL- PRESERVATIVE FREE
NDC 0409-1559-19
Rx only

Each mL contains 2.5 mg bupivacaine HCl, with NaCl to make isotonic. In Water for Injection, pH adjusted with NaOH or HCl.

Marcaine™ 0.25%
bupivacaine HCl injection, USP
75 mg/30 mL (2.5 mg/mL)
For INFILTRATION, NERVE BLOCK,
CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

Usual dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature]

Discard unused portion after initial use.

PAA160399

*****AA
DMMYYYYY

LOT
EXP

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA Hospira

PRINCIPAL DISPLAY PANEL - 75 mg/30 mL Vial Carton - 1559

30 mL

10 SINGLE-DOSE VIALS-PRESERVATIVE FREE

Rx only

NDC 0409-1559-30

Contains 10 of NDC 0409-1559-19

Marcaine™ 0.25%


bupivacaine HCl injection, USP

75 mg/30 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 50 mg/10 mL Vial Label - 1560

10 mL
 Single-dose Vial
 NDC 0409-1560-18
 Rx only


Preservative-Free

Marcaïne™ 0.5%
 bupivacaine HCl injection, USP
 50 mg/10 mL (5 mg/mL)

For NERVE BLOCK, CAUDAL
 and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 50 mg/10 mL Vial Tray - 1560

10 mL

Preservative-Free

10 Single-dose Vials

Rx only

NDC 0409-1560-10

Contains 10 of NDC 0409-1560-18

Marcaine™ 0.5%


bupivacaine HCl injection, USP

50 mg/10 mL (5 mg/mL)

For NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 150 mg/30 mL Vial Label - 1560

30 mL

SINGLE-DOSE VIAL- PRESERVATIVE FREE

NDC 0409-1560-19

Rx only

Marcaine™ 0.5%

bupivacaine HCl injection, USP

150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL,
and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL
ANESTHESIA

Hospira

FPO 2D 15 MIL
PAA160443

30 mL NDC 0409-1560-19
SINGLE-DOSE VIAL - PRESERVATIVE FREE Rx only

Marcaine™ 0.5%
bupivacaine HCl injection, USP
150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL,
and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

Hospira

Discard unused portion after initial use.

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

PAA 160443

LOT #####
EXP DMMYYYY

Each mL contains 5 mg bupivacaine HCl, with NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.
Usual dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.]

FPO DataBar Limited 10 MIL

PRINCIPAL DISPLAY PANEL - 150 mg/30 mL Vial Carton - 1560

30 mL
10 SINGLE-DOSE VIALS-PRESERVATIVE FREE
Rx only
NDC 0409-1560-29
Contains 10 of NDC 0409-1560-19

Marcaine™ 0.5%
bupivacaine HCl injection, USP
150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

Hospira

30 mL NDC 0409-1560-29
10 SINGLE-DOSE VIALS-PRESERVATIVE FREE Rx only
Contains 10 of NDC 0409-1560-19

Marcaine™ 0.5%
bupivacaine HCl injection, USP
150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

Hospira

Discard unused portion after initial use. Each mL contains 5 mg bupivacaine HCl, with NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.
Usual dosage: See insert. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F).
[See USP controlled room temperature.]

MADE IN SWEDEN
Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

30 mL NDC 0409-1560-29
10 SINGLE-DOSE VIALS-PRESERVATIVE FREE Rx only
Contains 10 of NDC 0409-1560-19

Marcaine™ 0.5%
bupivacaine HCl injection, USP
150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

Hospira

PRINCIPAL DISPLAY PANEL - 75 mg/10 mL Vial Label - 1582

10 mL
Single-dose Vial

NDC 0409-1582-18
Rx only


Preservative-Free

Marcaine™ 0.75%
bupivacaine HCl injection, USP
75 mg/10 mL (7.5 mg/mL)

EPIDURAL and RETROBULBAR ANESTHESIA USE

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 75 mg/10 mL Vial Tray - 1582

10 mL
10 Single-dose Vials

NDC 0409-1582-10
Contains 10 of NDC 0409-1582-18
Rx only

Preservative-Free

Marcaine™ 0.75%
bupivacaine HCl injection, USP
75 mg/10 mL (7.5 mg/mL)

EPIDURAL and RETROBULBAR ANESTHESIA USE

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 225 mg/30 mL Vial Label - 1582


30 mL
SINGLE-DOSE VIAL- PRESERVATIVE FREE

NDC 0409-1582-19
Rx only

Marcaïne™ 0.75%
bupivacaine HCl injection, USP
225 mg/30 mL (7.5 mg/mL)

for EPIDURAL and RETROBULBAR ANESTHESIA
NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 225 mg/30 mL Vial Carton - 1582

30 mL
10 SINGLE-DOSE VIALS-PRESERVATIVE FREE

NDC 0409-1582-29

Contains 10 of NDC 0409-1582-19
Rx only


Marcaine™ 0.75%
bupivacaine HCl injection, USP

225 mg/30 mL (7.5 mg/mL)

for EPIDURAL and RETROBULBAR ANESTHESIA

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Hospira



PRINCIPAL DISPLAY PANEL - 125 mg/50 mL Vial Label - 1587

50 mL Multiple-Dose vial

Marcaine™ 0.25%
bupivacaine HCl injection, USP

125 mg/50 mL
(2.5 mg/mL)

Each mL contains 2.5 mg bupivacaine HCl, with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

For INFILTRATION and NERVE BLOCK

50 mL Multiple-Dose vial

NDC 0409-1587-50

Rx only

Marcaine™ 0.25%

bupivacaine HCl injection, USP

125 mg/50 mL

(2.5 mg/mL)

Each mL contains 2.5 mg bupivacaine HCl, with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

For INFILTRATION and NERVE BLOCK

Not for caudal, epidural, or intrathecal anesthesia

Dosage: See package insert.

Store at 20°C to 25°C (68°F to 77°F).

[See USP Controlled Room Temperature.]

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA



(01) 00304091587502



PAA193030

#####AA

LOT

DDMMYY

EXP

PRINCIPAL DISPLAY PANEL - 125 mg/50 mL Vial Carton - 1587

NDC 0409-1587-50

Rx only

50 mL Multiple-Dose Vial

Marcaine™

0.25%

bupivacaine HCl injection, USP


125 mg/50 mL

(2.5 mg/mL)

For INFILTRATION
and NERVE BLOCK

Not for caudal, epidural,
or intrathecal anesthesia

Hospira



PRINCIPAL DISPLAY PANEL - 250 mg/50 mL Vial Label - 1610




50 mL Multiple-Dose Vial

Marcaine™ 0.5%
bupivacaine HCl injection, USP

250 mg/50 mL
(5 mg/mL)

Each mL contains 5 mg bupivacaine HCl,
with 1 mg methylparaben as antiseptic
preservative and NaCl to make isotonic,
in Water for Injection. pH adjusted with
NaOH or HCl.

for NERVE BLOCK

50 mL Multiple-Dose Vial	NDC 0409-1610-50	
Marcaine™ 0.5%	Rx only	
bupivacaine HCl injection, USP	Not for caudal, epidural, or intrathecal anesthesia	
250 mg/50 mL	Dosage: See insert. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]	
(5 mg/mL)		
Each mL contains 5 mg bupivacaine HCl, with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.		
for NERVE BLOCK		
	Distributed by Hospira, Inc. Lake Forest, IL 60045 USA	
		
	FPO - GS1 Databar Limited - 10 mil (01)00304091610507	
	 (01)00304091610507	
	PAA193033	
	#####AA	LOT
	#####YY	EXP

PRINCIPAL DISPLAY PANEL - 250 mg/50 mL Vial Carton - 1610

NDC 0409-1610-50
Rx only

50 mL Multiple-Dose Vial


Marcaine™
0.5%
bupivacaine HCl injection, USP

250 mg/50 mL
(5 mg/mL)

for NERVE BLOCK

Not for caudal, epidural,
or intrathecal anesthesia

Hospira



PRINCIPAL DISPLAY PANEL - 25 mg/10 mL Vial Label - 1746

Preservative-Free
10 mL Single-dose Vial


Marcaine™ 0.25%
25 mg/10 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride
and epinephrine injection, USP

For INFILTRATION, NERVE BLOCK,
CAUDAL and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia



PRINCIPAL DISPLAY PANEL - 25 mg/10 mL Vial Carton - 1746

10 mL 10 Single-dose Vials
Preservative-Free

NDC 0409-1746-10
Contains 10 of NDC 0409-1746-70
Rx only

Marcaine™ 0.25%
25 mg/10 mL (2.5 mg/mL)
with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP

For INFILTRATION, NERVE BLOCK, CAUDAL and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites

Hospira

FPO 20 15 MH
PAA160348



PAA160348

10 mL 10 Single-dose Vials
Preservative-Free

NDC 0409-1746-10
Contains 10 of NDC 0409-1746-70
Rx only

Marcaine™ 0.25%

25 mg/10 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP
For INFILTRATION, NERVE BLOCK, CAUDAL and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites



10 mL
Preservative-Free

NDC 0409-1746-10

Marcaine™ 0.25%

25 mg/10 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP

For INFILTRATION, NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites



WARNING: Contains Sulfites
Not for intrathecal anesthesia

bupivacaine hydrochloride and epinephrine injection, USP
For INFILTRATION, NERVE BLOCK, CAUDAL and EPIDURAL ANESTHESIA

with epinephrine 1:200,000 (as bitartrate)

25 mg/10 mL (2.5 mg/mL)

Marcaine™ 0.25%

Preservative-Free

NDC 0409-1746-10

10 mL 10 Single-dose Vials



LN *****
EXP 2M1MYY
SN *****

Contains epinephrine – do not autoclave.

Each mL contains 2.5 mg bupivacaine HCl and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, and 2 mg ascorbic acid as antioxidants, 0.0017 mL 60% sodium lactate buffer, 0.1 mg edetate calcium disodium as stabilizer, and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

Usual dosage: See Insert. **Protect from light.** This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Discard unused portion.

PAA160348

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA



FPOG1-128 10MH



(01) 10304091746104

PRINCIPAL DISPLAY PANEL - 75 mg/30 mL Vial Label - 1746

30 mL Single Dose Vial

NDC 0409-1746-71

Rx only

PRESERVATIVE-FREE

Marcaine™ 0.25%

75 mg/30 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP

for INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites

Hospira

30 mL Single Dose Vial NDC 0409-1746-71 Rx only

PRESERVATIVE-FREE

Marcaine™ 0.25%
75 mg/30 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP
for INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites

Each mL contains 2.5 mg bupivacaine HCl, and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothio glycerol, 2 mg ascorbic acid, 0.0017 mL 60% sodium lactate, 0.1 mg edetate calcium disodium, and NaCl to make isotonic pH adjusted with NaOH or HCl.

Usual dosage: See package insert.

Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]

Contains epinephrine-do not autoclave. Discard unused portion after initial use.

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

FPO 2D 15 Mil PAA160349

Lot: #####A
Exp: DMMYY

FPO - GS1 Databar Limited 10 Mil

PRINCIPAL DISPLAY PANEL - 75 mg/30 mL Vial Carton - 1746

30 mL 10 Single Dose Vials

NDC 0409-1746-30

Contains 10 of NDC 0409-1746-71

Rx only

PRESERVATIVE-FREE

Marcaine™ 0.25%

75 mg/30 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP

for INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites

Hospira



PAA160350

30 mL 10 Single Dose Vials

NDC 0409-1746-30
Contains 10 of NDC 0409-1746-71

Rx only

PRESERVATIVE-FREE

Marcaine™ 0.25%

75 mg/30 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP
for INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites



30 mL

NDC 0409-1746-30
Rx only

PRESERVATIVE-FREE

Marcaine™ 0.25%

75 mg/30 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP

for INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites



Warning: Contains Sulfites

Not for intrathecal anesthesia

bupivacaine hydrochloride and epinephrine injection, USP
for INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

with epinephrine 1:200,000 (as bitartrate)


75 mg/30 mL (2.5 mg/mL)

Marcaine™ 0.25%

PRESERVATIVE-FREE

30 mL 10 Single Dose Vials

NDC 0409-1746-30
Rx only



Contains epinephrine – do not autoclave.

Discard unused portion after initial use.

Each mL contains 2.5 mg bupivacaine HCl and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, and 2 mg ascorbic acid as antioxidants, 0.0017 mL 60% sodium lactate buffer, 0.1 mg edetate calcium disodium as stabilizer, and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl. **Usual dosage:** See package insert.

Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.]



PAA160350

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA



CU174630204091746302



PRINCIPAL DISPLAY PANEL - 50 mg/10 mL Vial Label - 1749

10 mL Single-dose Vial
Preservative-Free

Marcaine™ 0.5%
50 mg/10 mL (5 mg/mL)
with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride
and epinephrine injection, USP

For NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites

Hospira

10 mL Single-dose Vial Preservative-Free NDC 0409-1749-70 Rx only

Marcaine™ 0.5%
50 mg/10 mL (5 mg/mL)
with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride
and epinephrine injection, USP

**For NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA**

Not for intrathecal anesthesia

WARNING: Contains Sulfites

Each mL contains 5 mg bupivacaine HCl and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, 2 mg ascorbic acid, 0.0017 mL 60% sodium lactate, 0.1 mg edetate calcium disodium, and NaCl to make isotonic. pH adjusted with NaOH or HCl. **Usual dosage:** See insert. **Protect from light.** This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.] Discard unused portion.

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

Contains epinephrine – do not autoclave.

QR Code

Barcode

PAA160351

LOT/EXP

#####AA
DMMYYY

FPO - 2D 15 Mil
PAA160351

FPO - GS1 Databar Limited 7.5 Mil

PRINCIPAL DISPLAY PANEL - 50 mg/10 mL Vial Carton - 1749

10 mL 10 Single-dose Vials

NDC 0409-1749-10
Contains 10 of NDC 0409-1749-70
Rx only

Preservative-Free

Marcaine™ 0.5%
50 mg/10 mL (5 mg/mL)
with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP

For NERVE BLOCK, CAUDAL and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites

Hospira



PAA160352

10 mL 10 Single-dose Vials

NDC 0409-1749-10
Contains 10 of NDC 0409-1749-70

Rx only

Preservative-Free

Marcaine™ 0.5%

50 mg/10 mL (5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP
For NERVE BLOCK, CAUDAL and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites



Preservative-Free

Marcaine™ 0.5%

50 mg/10 mL (5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP
For NERVE BLOCK, CAUDAL and
EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

WARNING: Contains Sulfites



WARNING: Contains Sulfites

Not for intrathecal anesthesia

bupivacaine hydrochloride and epinephrine injection, USP
For NERVE BLOCK, CAUDAL and EPIDURAL ANESTHESIA

with epinephrine 1:200,000 (as bitartrate)

50 mg/10 mL (5 mg/mL)

Marcaine™ 0.5%

Preservative-Free

LOT: *****
EXP: DMMYY
GSI Data Mark

GM (013060974905)

Contains epinephrine – do not autoclave.

Each mL contains 5 mg bupivacaine HCl and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, and 2 mg ascorbic acid as antioxidants, 0.0017 mL 60% sodium lactate buffer, 0.1 mg edetate calcium disodium as stabilizer, and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

Usual dosage: See insert. Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Discard unused portion.

PAA160352


Distributed by Hospira, Inc., Lake Forest, IL 60045 USA



FGCSI-128 10 ML



(01)10304091749105



PRINCIPAL DISPLAY PANEL - 150 mg/30 mL Vial Label - 1749

30 mL
Single Dose Vial

NDC 0409-1749-71
Rx only

PRESERVATIVE-FREE

Marcaine™ 0.5%
150 mg/30 mL (5 mg/mL)
with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP

for NERVE BLOCK, CAUDAL, and EPIDURAL
ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites

Hospira

FPO - 2D 15 MII
PAA175213

30 mL
Single Dose Vial
PRESERVATIVE-FREE

NDC 0409-1749-71
Rx only

Marcaine™ 0.5%
150 mg/30 mL (5 mg/mL)
with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine Injection, USP
for NERVE BLOCK, CAUDAL, and EPIDURAL
ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites

Each mL contains 5 mg bupivacaine HCl and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothoglycerol, 2 mg ascorbic acid, 0.0017 mL 60% sodium lactate, 0.1 mg edetate calcium disodium, and NaCl to make isotonic pH adjusted with NaOH or HCl.

Usual dosage: See package insert. Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]

Contains epinephrine - do not autoclave
Discard unused portion after initial use

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

(01)00304091749719

PAA175213

#####AA
DMMYYYY

LOT
EXP

FPO - GS1 dataBar I limited 10 mil

PRINCIPAL DISPLAY PANEL - 150 mg/30 mL Vial Carton - 1749

30 mL 10 Single Dose Vials

NDC 0409-1749-29

Contains 10 of NDC 0409-1749-71

Rx only

PRESERVATIVE-FREE

Marcaine™ 0.5%

150 mg/30 mL (5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP
for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites

Hospira



PAA160354

30 mL 10 Single Dose Vials

PRESERVATIVE-FREE

Marcaine™ 0.5%

150 mg/30 mL (5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and epinephrine injection, USP
for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia **Warning: Contains Sulfites**



NDC 0409-1749-29
Contains 10 of NDC 0409-1749-71

Rx only

30 mL

NDC 0409-1749-29

PRESERVATIVE-FREE

Marcaine™ 0.5%

150 mg/30 mL (5 mg/mL)

**with epinephrine
1:200,000 (as bitartrate)**

bupivacaine hydrochloride and
epinephrine injection, USP
for NERVE BLOCK, CAUDAL,
and EPIDURAL ANESTHESIA

Not for intrathecal anesthesia

Warning: Contains Sulfites



Not for intrathecal anesthesia **Warning: Contains Sulfites**
bupivacaine hydrochloride and epinephrine injection, USP
for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

with epinephrine 1:200,000 (as bitartrate)

150 mg/30 mL (5 mg/mL)


Marcaine™ 0.5%

PRESERVATIVE-FREE

Rx only

NDC 0409-1749-29

30 mL 10 Single Dose Vials



Contains epinephrine – do not autoclave.
Discard unused portion after initial use.

Each mL contains 5 mg bupivacaine HCl and 0.0091 mg epinephrine bitartrate, with 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, and 2 mg ascorbic acid as antioxidants, 0.0017 mL 60% sodium lactate buffer, 0.1 mg edetate calcium disodium as stabilizer, and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

Usual dosage: See package insert.

Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.]

PAA160354




PAA160354

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA



0171 1030409174929



PRINCIPAL DISPLAY PANEL - 125 mg/50 mL Vial Label - 1752

50 mL Multiple-Dose Vial

NDC 0409-1752-50

Rx only

Marcaine™ 0.25%

125 mg/50 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP
for INFILTRATION and NERVE BLOCK

Not for caudal, epidural, or intrathecal anesthesia

Contains epinephrine—do not autoclave

Warning: Contains Sulfites

FPO - 2D - 15 mil
PAA174206

50 mL Multiple-Dose Vial NDC 0409-1752-50
Rx only

Marcaine™ 0.25%
125 mg/50 mL (2.5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP
for INFILTRATION and NERVE BLOCK

Not for caudal, epidural, or intrathecal anesthesia

Contains epinephrine—do not autoclave
Warning: Contains Sulfites

Each mL contains 2.5 mg bupivacaine HCl, and 0.0091 mg epinephrine bitartrate, with 1 mg methylparaben as preservative, 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, 2 mg ascorbic acid, 0.0017 mL 60% sodium lactate, 0.1 mg edetate calcium disodium, and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl. Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]
Dosage: See package insert.

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

FPO - GS1 Databar Limited - 10 mil

PAA174206

LOT ###A
EXP DMMYYY

PRINCIPAL DISPLAY PANEL - 125 mg/50 mL Vial Carton - 1752

NDC 0409-1752-50

Rx only

50 mL Multiple-Dose Vial

Marcaine™

0.25%

125 mg/50 mL

(2.5 mg/mL)

with epinephrine 1:200,000

(as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP


for INFILTRATION
and NERVE BLOCK

Not for caudal, epidural,
or intrathecal anesthesia

Contains epinephrine-
do not autoclave

Warning: Contains Sulfites

Hospira



PRINCIPAL DISPLAY PANEL - 250 mg/50 mL Vial Label - 1755

50 mL Multiple-Dose Vial

NDC 0409-1755-50

Rx only

Marcaine™ 0.5%

250 mg/50 mL (5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP
for NERVE BLOCK

Not for caudal, epidural, or intrathecal anesthesia

Contains epinephrine—do not autoclave

Warning: Contains Sulfites

50 mL Multiple-Dose Vial NDC 0409-1755-50 Rx only

Marcaine™ 0.5%
250 mg/50 mL (5 mg/mL)

with epinephrine 1:200,000 (as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP
for NERVE BLOCK

Not for caudal, epidural, or intrathecal anesthesia

Contains epinephrine—do not autoclave
Warning: Contains Sulfites

Each mL contains 5 mg bupivacaine HCl, and 0.0091 mg epinephrine bitartrate, with 1 mg methylparaben as preservative, 0.5 mg sodium metabisulfite, 0.001 mL monothioglycerol, 2 mg ascorbic acid, 0.0017 mL 60% sodium lactate, 0.1 mg edetate calcium disodium, and NaCl to make isotonic. pH adjusted with NaOH or HCl. Protect from light. This product is clear and colorless. Do not use the solution if it is discolored or if it contains a precipitate.
Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]
Usual dosage: See package insert.

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

FPO - 2D - 15 mil
PAA174207

FPO - GS1 DataBar Limited 10 mil

LOT ###AA
EXP DMMYYY

Hospira

PRINCIPAL DISPLAY PANEL - 250 mg/50 mL Vial Carton - 1755

NDC 0409-1755-50

Rx only

50 mL Multiple-Dose Vial

Marcaine™
0.5%

250 mg/50 mL
(5 mg/mL)

with epinephrine 1:200,000
(as bitartrate)

bupivacaine hydrochloride and
epinephrine injection, USP

for NERVE BLOCK


Not for caudal, epidural, or
intrathecal anesthesia

Contains epinephrine—

do not autoclave

Warning: Contains Sulfites

Hospira



PRINCIPAL DISPLAY PANEL - 25 mg/10 mL Vial Label - 2510


10 mL
Preservative-Free
Single-dose Vial
NDC 0409-2510-01
Rx only

Marcaine™ 0.25%
bupivacaine HCl injection, USP
25 mg/10 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA.

AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/F/R
Dist. by Hospira, Inc., Lake Forest, IL 60045 USA
Hospira

10 mL Single-dose Vial NDC 0409-2510-01
Preservative-Free Rx only
Marcaine™ 0.25%
bupivacaine HCl injection, USP
25 mg/10 mL (2.5 mg/mL)
For INFILTRATION, NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA.
AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/F/R
Dist. by Hospira, Inc., Lake Forest, IL 60045 USA 

Each mL contains 2.5 mg bupivacaine HCl with NaCl to make isotonic, in Water for Injection, pH adjusted with NaOH or HCl.
Usual Dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]
Discard unused portion.

PAA223300

PRINCIPAL DISPLAY PANEL - 25 mg/10 mL Vial Tray - 2510

10 mL
Preservative-Free
25 Single-dose Vials

NDC 0409-2510-25
Contains 25 of NDC 0409-2510-01
Rx only

Marcaine™ 0.25%
bupivacaine HCl injection, USP
25 mg/10 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

10 mL 25 Single-dose Vials NDC 0409-2510-25
Preservative-Free Contains 25 of NDC 0409-2510-01
Rx only

Marcaine™ 0.25%
bupivacaine HCl injection, USP

25 mg/10 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA



(01)00304092510257



AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/F/R
Distributed by Hospira, Inc., Lake Forest, IL 60045 USA
MADE IN SWEDEN

Discard unused portion.
Each mL contains 2.5 mg bupivacaine HCl, with NaCl to make isotonic, in Water for Injection, pH adjusted with NaOH or HCl.
Usual Dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature.]

PA A223321

10 mL 25 Single-dose Vials
Preservative-Free
Marcaine™ 0.25%
bupivacaine HCl injection, USP
25 mg/10 mL (2.5 mg/mL)
For INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

NDC 0409-2510-25



PRINCIPAL DISPLAY PANEL - 75 mg/30 mL Vial Label - 7535

30 mL
SINGLE-DOSE VIAL- PRESERVATIVE FREE

NDC 0409-7535-01
Rx only

Marcaine™ 0.25%
bupivacaine HCl injection, USP

75 mg/30 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK,
CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA
Hospira

Discard unused portion
after initial use.


AP/DRUGS/08/2013

M.L.No. 08/VP/AP/2013/F/R

30 mL
SINGLE-DOSE VIAL - PRESERVATIVE FREE
NDC 0409-7535-01
Rx only

Marcaïne™ 0.25%
bupivacaine HCl injection, USP
75 mg/30 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK,
CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA 

Usual dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.]

Discard unused portion
after initial use.

M.L.No. 08/VP/AP/2013/F/R
AP/DRUGS/08/2013

PAA223322

PRINCIPAL DISPLAY PANEL - 75 mg/30 mL Vial Tray - 7535

30 mL

25 SINGLE-DOSE VIALS-PRESERVATIVE FREE

NDC 0409-7535-25

Contains 25 of NDC 0409-7535-01

Rx only


Marcaïne™ 0.25%

bupivacaine HCl injection, USP

75 mg/30 mL (2.5 mg/mL)

For INFILTRATION, NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA



PRINCIPAL DISPLAY PANEL - 125 mg/50 mL Vial Label - 1250

50 mL Multiple-Dose Vial

NDC 0409-1250-01

Rx only

Marcaine™ 0.25%
bupivacaine HCl injection, USP

125 mg/50 mL
(2.5 mg/mL)

Each mL contains 2.5 mg bupivacaine

HCl, with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

For INFILTRATION and NERVE BLOCK

Not for caudal, epidural, or intrathecal anesthesia



Dosage: See package insert.

Store at 20°C to 25°C (68°F to 77°F).[See USP Controlled Room Temperature.]

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA
Hospira

AP/DRUGS/08/2013

M.L.No.08/VP/AP/2013/F/R

<p>50 mL Multiple-Dose vial</p> <p>Marcaine™ 0.25% bupivacaine HCl injection, USP 125 mg/50 mL (2.5 mg/mL)</p> <p>Each mL contains 2.5 mg bupivacaine HCl, with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.</p> <p>For INFILTRATION and NERVE BLOCK</p>	<p>NDC 0409-1250-01 Rx only</p> <p>Not for caudal, epidural, or intrathecal anesthesia</p> <p>Dosage: See package insert.</p> <p>Store at 20°C to 25°C (68°F to 77°F).[See USP Controlled Room Temperature.]</p> <p>Distributed by Hospira, Inc. Lake Forest, IL 60045 USA AP/DRUGS/08/2013 M.L.No.08/VP/AP/2013/F/R</p> 	 <p>(01)10304091250014</p> <p>PAA223324</p>
---	--	---

PRINCIPAL DISPLAY PANEL - 125 mg/50 mL Vial Tray - 1250

50 mL Multiple-Dose Vial

Rx only

NDC 0409-1250-25

Contains 25 of NDC 0409-1250-01

Marcaine™ 0.25%
bupivacaine HCl injection, USP

125 mg/50 mL (2.5 mg/mL)

For INFILTRATION

and NERVE BLOCK

Not for caudal, epidural,
or intrathecal anesthesia

Hospira

50 mL Multiple-Dose Vial

Rx only NDC 0409-1250-25
Contains 25 of NDC 0409-1250-01

Marcaine™ 0.25%
bupivacaine HCl injection, USP
125 mg/50 mL (2.5 mg/mL)

For INFILTRATION
and NERVE BLOCK
Not for caudal, epidural,
or intrathecal anesthesia

Hospira

(01)00304091250253

MADE IN SWEDEN
Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/FF/R

Each mL contains 2.5 mg bupivacaine HCl, with 1 mg methylparaben as
antiseptic preservative and NaCl to make isotonic in Water for Injection,
pH adjusted with NaOH or HCl.
Dosage: See package insert.
This product is clear and colorless. Do not use the solution if it is discolored
or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F); excursions
permitted between 15°C to 30°C (59°F to 86°F). [See USP Controlled Room
Temperature.]

Hospira

50 mL Multiple-Dose Vial

Rx only NDC 0409-1250-25

For INFILTRATION
and NERVE BLOCK
Not for caudal, epidural,
or intrathecal anesthesia

Hospira

125 mg/50 mL (2.5 mg/mL)
bupivacaine HCl injection, USP
Marcaine™ 0.25%

PA A 22 3 325

PRINCIPAL DISPLAY PANEL - 50 mg/10 mL Vial Label - 5010

10 mL
Preservative-Free

Single-dose Vial

NDC 0409-5010-01

Rx only

Marcaine™ 0.5%
bupivacaine HCl injection, USP
50 mg/10 mL (5 mg/mL)

For NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA.

AP/DRUGS/08/2013

M.L.No.08/VP/AP/2013/F/R

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA
Hospira

10 mL Single-dose Vial NDC 0409-5010-01
Preservative-Free Rx only



Marcaine™ 0.5%
bupivacaine HCl injection, USP
50 mg/10 mL (5 mg/mL)

For NERVE BLOCK, CAUDAL
and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA.

AP/DRUGS/08/2013
M.L.No.08/VP/AP/2013/F/R
Dist. by Hospira, Inc., Lake Forest, IL 60045 USA

Each mL contains 5 mg bupivacaine HCl, with NaCl to make isotonic, in Water for Injection, pH adjusted with NaOH or HCl. Usual Dosage: See insert. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.] Discard unused portion.

PAA223294



PRINCIPAL DISPLAY PANEL - 50 mg/10 mL Vial Tray - 5010

10 mL
Preservative-Free

25 Single-dose Vials

NDC 0409-5010-25
Contains 25 of NDC 0409-5010-01
Rx only

Marcaine™ 0.5%
bupivacaine HCl injection, USP

50 mg/10 mL (5 mg/mL)

For NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

10 mL
Preservative-Free
Marcaine™ 0.5%
bupivacaine HCl injection, USP
50 mg/10 mL (5 mg/mL)
For NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

25 Single-dose Vials
Contains 25 of NDC 0409-5010-01

NDC 0409-5010-25
NDC 0409-5010-01

Rx only



(01)00304095010259



Discard unused portion.
Each mL contains 5 mg bupivacaine HCl, with NaCl to make isotonic
in Water for Injection, pH adjusted with NaOH or HCl.
Usual Dosage: See in sert.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between
15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature.]

AP/DRUGS/08/2013
ML No. 08/AP/AP/2013/F/R
MADE IN SWEDEN
Distributed by Hospira, Inc., Lake Forest, IL 60045 USA



5 AA 223295

10 mL
Preservative-Free
25 Single-dose Vials
NDC 0409-5010-25
Marcaine™ 0.5%
bupivacaine HCl injection, USP
50 mg/10 mL (5 mg/mL)
For NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

PRINCIPAL DISPLAY PANEL - 150 mg/30 mL Vial Label - 1530

30 mL
SINGLE-DOSE VIAL- PRESERVATIVE FREE

NDC 0409-1530-01

Rx only

Marcaine™ 0.5%
bupivacaine HCl injection, USP

150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL,

and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA

AP/DRUGS/08/2013

M.L.No 08/VP/AP/2013/F/R

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA


Hospira

Discard unused portion after
initial use.

30 mL
SINGLE-DOSE VIAL- PRESERVATIVE FREE

NDC 0409-1530-01
Rx only

Marcaine™ 0.5%
bupivacaine HCl injection, USP
150 mg/30 mL (5 mg/mL)
for NERVE BLOCK, CAUDAL,
and EPIDURAL ANESTHESIA
NOT FOR INTRATHECAL ANESTHESIA

AP/DRUGS/08/2013
M.L.No 08/VP/AP/2013/F/R
Dist. by Hospira, Inc., Lake Forest, IL 60045 USA 

Each mL contains 5 mg bupivacaine HCl,
with NaCl to make isotonic, in Water for
Injection; pH adjusted with NaOH or HCl.
Usual dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.]

Discard unused portion after
initial use.

PAA223296

PRINCIPAL DISPLAY PANEL - 150 mg/30 mL Vial Tray - 1530

30 mL

25 Single-dose Vials – Preservative Free

NDC 0409-1530-25

Contains 25 of NDC 0409-1530-01

Rx only


Marcaine™ 0.5%

bupivacaine HCl injection, USP

150 mg/30 mL (5 mg/mL)

for NERVE BLOCK, CAUDAL, and EPIDURAL ANESTHESIA

NOT FOR INTRATHECAL ANESTHESIA



PRINCIPAL DISPLAY PANEL - 250 mg/50 mL Vial Label - 0525

50 mL Multiple-Dose Vial

NDC 0409-0525-01

Rx only

Marcaine™ 0.5%
bupivacaine HCl injection, USP

250 mg/50 mL
(5 mg/mL)

Each mL contains 5 mg bupivacaine HCl,

with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.

for NERVE BLOCK

Distributed by Hospira, Inc. Lake Forest, IL 60045 USA

Not for caudal, epidural, or intrathecal anesthesia

Dosage: See insert.

Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]

AP/DRUGS/08/2013

M.L.No. 08/VP/AP/2013/F/R

Hospira

50 mL Multiple-Dose Vial
Marcaine™ 0.5%
bupivacaine HCl injection, USP
250 mg/50 mL
(5 mg/mL)

Each mL contains 5 mg bupivacaine HCl, with 1 mg methylparaben as antiseptic preservative and NaCl to make isotonic, in Water for Injection. pH adjusted with NaOH or HCl.
for NERVE BLOCK

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

NDC 0409-0525-01
Rx only

Not for caudal, epidural, or intrathecal anesthesia

Dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]

AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/F/R

(01)10304090525014
PAA223298

PRINCIPAL DISPLAY PANEL - 250 mg/50 mL Vial Tray - 0525

50 mL Multiple-Dose Vial

Rx only

NDC 0409-0525-25

Contains 25 of NDC 0409-0525-01

Marcaine™ 0.5%

bupivacaine HCl injection, USP

250 mg/50 mL (5 mg/mL)

For NERVE BLOCK

Not for caudal, epidural,
or intrathecal anesthesia

Hospira

50 mL Multiple-Dose Vial

Rx only NDC 0409-0525-25
Contains 25 of NDC 0409-0525-01

Marcaine™ 0.5%
bupivacaine HCl injection, USP
250 mg/50 mL (5 mg/mL)

For NERVE BLOCK
Not for caudal, epidural,
or intrathecal anesthesia

Hospira

(01)003040V0525253

MADE IN SWEDEN
Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013.F/R

Each mL contains 5 mg bupivacaine HCl, with 1 mg methylparaben as
antiseptic preservative and NaCl to make isotonic, in Water for Injection,
pH adjusted with NaOH or HCl.
Dosage: See package insert.
This product is clear and colorless. Do not use the solution if it is discolored
or if it contains a precipitate. Store at 20°C to 25°C (68°F to 77°F); excursions
permitted between 15°C to 30°C (59°F to 86°F). [See USP Controlled Room
Temperature.]

Hospira

50 mL Multiple-Dose Vial

Rx only NDC 0409-0525-25

Marcaine™ 0.5%
bupivacaine HCl injection, USP
250 mg/50 mL (5 mg/mL)

For NERVE BLOCK
Not for caudal, epidural,
or intrathecal anesthesia

Hospira

PA223299

PRINCIPAL DISPLAY PANEL - 75 mg/10 mL Vial Label - 7510

10 mL
Single-dose Vial
Preservative-Free

NDC 0409-7510-01

Rx only

Marcaine™ 0.75%

bupivacaine HCl injection, USP

75 mg/10 mL (7.5 mg/mL)

EPIDURAL and RETROBULBAR ANESTHESIA USE

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

AP/DRUGS/08/2013

M.L.No. 08/VP/AP/2013/F/R

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA

Hospira

Discard unused portion.

10 mL
Single-dose Vial
Preservative-Free

Marcaine™ 0.75%
bupivacaine HCl injection, USP
75 mg/10 mL (7.5 mg/mL)
EPIDURAL and RETROBULBAR ANESTHESIA USE
NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Each mL contains 7.5 mg bupivacaine HCl, with NaCl to make isotonic, in Water for Injection, pH adjusted with NaOH or HCl.

Usual Dosage: See insert.
Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.]

NDC 0409-7510-01
Rx only

AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/F/R
Dist. by Hospira, Inc., Lake Forest, IL 60045 USA

Discard unused portion.

PAA223326

PRINCIPAL DISPLAY PANEL - 75 mg/10 mL Vial Tray - 7510

10 mL

Preservative-Free

25 Single-dose Vials

NDC 0409-7510-25

Contains 25 of NDC 0409-7510-01

Rx only


Marcaine™ 0.75%

bupivacaine HCl injection, USP

75 mg/10 mL (7.5 mg/mL)

EPIDURAL and RETROBULBAR ANESTHESIA USE

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA



PRINCIPAL DISPLAY PANEL - 225 mg/30 mL Vial Label - 2253

30 mL
SINGLE-DOSE VIAL- PRESERVATIVE FREE

NDC 0409-2253-01
Rx only

Marcaine™ 0.75%
bupivacaine HCl injection, USP

225 mg/30 mL (7.5 mg/mL)

for EPIDURAL and RETROBULBAR ANESTHESIA

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA

AP/DRUGS/08/2013

M.L.No. 08/VP/AP/2013/F/R

Hospira

Discard unused portion after
initial use.

30 mL NDC 0409-2253-01
SINGLE-DOSE VIAL - PRESERVATIVE FREE Rx only

Marcaine™ 0.75%
bupivacaine HCl injection, USP
225 mg/30 mL (7.5 mg/mL)

for EPIDURAL and RETROBULBAR ANESTHESIA
NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA
AP/DRUGS/08/2013
M.L.No. 08/VP/AP/2013/F/R

Each mL contains 7.5 mg bupivacaine HCl, with NaCl to make isotonic, in Water for Injection, pH adjusted with NaOH or HCl.

Usual dosage: See insert.

Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature.]

Discard unused portion after
initial use.

PAA223328

PRINCIPAL DISPLAY PANEL - 225 mg/30 mL Vial Tray - 2253

30 mL

25 SINGLE-DOSE VIALS-PRESERVATIVE FREE

NDC 0409-2253-25

Contains 25 of NDC 0409-2253-01

Rx only


Marcaine™ 0.75%

bupivacaine HCl injection, USP

225 mg/30 mL (7.5 mg/mL)

for EPIDURAL and RETROBULBAR ANESTHESIA

NOT FOR OBSTETRIC OR INTRATHECAL ANESTHESIA



MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1559
Route of Administration	EPIDURAL, INFILTRATION, PERINEURAL, INTRACAUDAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)		BUPIVACAINE HYDROCHLORIDE ANHYDROUS	2.5 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1559-10	10 in 1 TRAY	08/23/2005	
1	NDC:0409-1559-18	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-1559-30	10 in 1 CARTON	09/08/2005	
2	NDC:0409-1559-19	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	08/23/2005	

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1560
Route of Administration	EPIDURAL, PERINEURAL, INTRACAUDAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	5 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1560-10	10 in 1 TRAY	08/31/2005	
1	NDC:0409-1560-18	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-1560-29	10 in 1 CARTON	08/16/2005	
2	NDC:0409-1560-19	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	08/16/2005	

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1582
Route of Administration	EPIDURAL, RETROBULBAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	7.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1582-10	10 in 1 TRAY	07/27/2005	
1	NDC:0409-1582-18	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-1582-29	10 in 1 CARTON	08/08/2005	
2	NDC:0409-1582-19	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	07/27/2005	

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1587
Route of Administration	INFILTRATION, PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	2.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1587-50	1 in 1 CARTON	01/17/2006	
1		50 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	01/17/2006	

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1610
Route of Administration	PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W471Q8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1610-50	1 in 1 CARTON	11/30/2005	
1		50 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	11/30/2005	

MARCAINE WITH EPINEPHRINE

bupivacaine hydrochloride and epinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1746
Route of Administration	INFILTRATION, EPIDURAL, INTRACAUDAL, PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	2.5 mg in 1 mL
EPINEPHRINE BITARTRATE (UNII: 3OQ7KI53AK) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE BITARTRATE	0.0091 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM LACTATE (UNII: TU7HW0W0QT)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	
WATER (UNII: 059QF0KO0R)	
MONOTHIOGLYCEROL (UNII: AAO1P0WSXJ)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1746-10	10 in 1 CARTON	12/09/2005	
1	NDC:0409-1746-70	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-1746-30	10 in 1 CARTON	09/12/2005	
2	NDC:0409-1746-71	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	09/12/2005	

MARCAINE WITH EPINEPHRINE

bupivacaine hydrochloride and epinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1749
Route of Administration	EPIDURAL, INTRACAUDAL, PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	5 mg in 1 mL
EPINEPHRINE BITARTRATE (UNII: 30Q7KI53AK) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE BITARTRATE	0.0091 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM LACTATE (UNII: TU7HW0W0QT)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	
WATER (UNII: 059QF0KO0R)	
MONOTHIOGLYCEROL (UNII: AAO1P0WSXJ)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1749-10	10 in 1 CARTON	09/22/2005	
1	NDC:0409-1749-70	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-1749-29	10 in 1 CARTON	09/12/2005	
2	NDC:0409-1749-71	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	09/12/2005	

MARCAINE WITH EPINEPHRINE

bupivacaine hydrochloride and epinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1752
Route of Administration	INFILTRATION, PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	2.5 mg in 1 mL
EPINEPHRINE BITARTRATE (UNII: 30Q7KI53AK) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE BITARTRATE	0.0091 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM LACTATE (UNII: TU7HW0W0QT)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	
MONOTHIOGLYCEROL (UNII: AAO1P0WSXJ)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1752-50	1 in 1 CARTON	02/06/2006	
1		50 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	02/06/2006	

MARCAINE WITH EPINEPHRINE

bupivacaine hydrochloride and epinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1755
Route of Administration	PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	5 mg in 1 mL
EPINEPHRINE BITARTRATE (UNII: 30Q7KI53AK) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE BITARTRATE	0.0091 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM LACTATE (UNII: TU7HW0W0QT)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	
MONOTHIOGLYCEROL (UNII: AAO1P0WSXJ)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1755-50	1 in 1 CARTON	03/22/2006	
1		50 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	03/22/2006	

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-2510
Route of Administration	EPIDURAL, INFILTRATION, PERINEURAL, INTRACAUDAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	25 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-2510-25	25 in 1 TRAY	12/09/2024	12/09/2024
1	NDC:0409-2510-01	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-7535
Route of Administration	EPIDURAL, INFILTRATION, PERINEURAL, INTRACAUDAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	75 mg in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

SODIUM HYDROXIDE (UNII: 55X04QC32I)

HYDROCHLORIC ACID (UNII: QTT17582CB)

WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-7535-25	25 in 1 TRAY	12/09/2024	12/09/2024
1	NDC:0409-7535-01	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1250
Route of Administration	INFILTRATION, PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	125 mg in 50 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1250-25	25 in 1 TRAY	12/09/2024	12/09/2024

1	NDC:0409-1250-01	50 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		
----------	------------------	--	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-5010
Route of Administration	EPIDURAL, PERINEURAL, INTRACAUDAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	50 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-5010-25	25 in 1 TRAY	12/09/2024	12/09/2024
1	NDC:0409-5010-01	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-1530
Route of Administration	EPIDURAL, PERINEURAL, INTRACAUDAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	150 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-1530-25	25 in 1 TRAY	12/09/2024	12/09/2024
1	NDC:0409-1530-01	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-0525
Route of Administration	PERINEURAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)		BUPIVACAINE HYDROCHLORIDE ANHYDROUS	250 mg in 50 mL	
Inactive Ingredients				
Ingredient Name		Strength		
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
HYDROCHLORIC ACID (UNII: QTT17582CB)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
WATER (UNII: 059QF0K00R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-0525-25	25 in 1 TRAY	12/09/2024	
1	NDC:0409-0525-01	50 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA016964	12/09/2024		

MARCAINE			
bupivacaine hydrochloride injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-7510
Route of Administration	EPIDURAL, RETROBULBAR		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)		BUPIVACAINE HYDROCHLORIDE ANHYDROUS	75 mg in 10 mL
Inactive Ingredients			
Ingredient Name		Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X)			
SODIUM HYDROXIDE (UNII: 55X04QC32I)			
HYDROCHLORIC ACID (UNII: QTT17582CB)			

WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-7510-25	25 in 1 TRAY	12/09/2024	12/09/2024
1	NDC:0409-7510-01	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

MARCAINE

bupivacaine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-2253
Route of Administration	EPIDURAL, RETROBULBAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUPIVACAINE HYDROCHLORIDE (UNII: 7TQO7W3VT8) (BUPIVACAINE - UNII:Y8335394RO)	BUPIVACAINE HYDROCHLORIDE ANHYDROUS	225 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-2253-25	25 in 1 TRAY	12/09/2024	12/09/2024
1	NDC:0409-2253-01	30 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016964	12/09/2024	12/09/2024

Labeler - Hospira, Inc. (141588017)

Establishment

Name	Address	ID/FEI	Business Operations
Hospira, Inc.		030606222	ANALYSIS(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-1746, 0409-1749, 0409-1752, 0409-1755) , MANUFACTURE(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-1746, 0409-1749, 0409-1752, 0409-1755) , PACK(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-1746, 0409-1749, 0409-1752, 0409-1755) , LABEL(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-1746, 0409-1749, 0409-1752, 0409-1755)

Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Healthcare India Private Limited		860037912	ANALYSIS(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-2510, 0409-7535, 0409-1250, 0409-5010, 0409-1530, 0409-0525, 0409-7510, 0409-2253) , MANUFACTURE(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-2510, 0409-7535, 0409-1250, 0409-5010, 0409-1530, 0409-0525, 0409-7510, 0409-2253) , PACK(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-2510, 0409-7535, 0409-1250, 0409-5010, 0409-1530, 0409-0525, 0409-7510, 0409-2253) , LABEL(0409-1559, 0409-1560, 0409-1582, 0409-1587, 0409-1610, 0409-2510, 0409-7535, 0409-1250, 0409-5010, 0409-1530, 0409-0525, 0409-7510, 0409-2253)

Revised: 12/2024

Hospira, Inc.